Downloaded from UvA-DARE, the institutional repository of the University of Amsterdam (UvA) http://dare.uva.nl/document/155519

File ID155519FilenameChapter 7: General discussion and summary

SOURCE (OI	R PART OF THE FOLLOWING SOURCE):
Туре	Dissertation
Title	Small GTPases : emerging targets in rheumatoid arthritis
Author	J.R. Ferreira de Abreu
Faculty	Faculty of Medicine
Year	2009
Pages	198

FULL BIBLIOGRAPHIC DETAILS: http://dare.uva.nl/record/323968

Copyright

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use.

UvA-DARE is a service provided by the library of the University of Amsterdam (http://dare.uva.nl)

7

GENERAL DISCUSSION AND SUMMARY

Rheumatoid arthritis (RA) is an auto-immune disease for which no cure is yet available. Recent insights into the pro-inflammatory cytokines and cells present in the inflamed joint, and their contributions to pathology in RA, have led to the development of biological therapies aimed at their neutralization. Anti-TNF- α blockade for example, is one of the most widely used and successful biological therapies¹. There is a large group of patients however, that don't respond sufficiently to treatment. Also, therapies with biologicals can have serious side effects as the immune system of treated patients become less efficient in clearing pathogens and opportunistic infections. Therefore, there is still the need for the development of new and more specific therapies.

Cellular activation, proliferation, survival and migration are tightly regulated at the intracellular level by Ras-superfamily small GTPases. Small GTPases are important modulators of cellular responses and crucial mediators of effective immune responses, and initial evidence has suggested these enzymes may contribute to pathology in RA. Ras family GTPase effector pathways such as MAP kinases, PI3K and NF-KB are highly activated in RA synovial tissue (ST)²⁻⁴. Moreover, strategies that block Ras function are able to suppress fibroblast-like synoviocyte (FLS) proliferation, invasiveness and pro-inflammatory cytokine production in vitro, and exert protective effects in animal models of arthritis in vivo⁵⁻⁷. Similarly, blockage of Rho family small GTPases reduces FLS proliferation, invasiveness, JNK activation and IL-6 production in vitro^{8,9}. Finally, inactivation of the Ras-related small GTPase Rap1 in synovial T cells is associated with increased intracellular oxidative stress¹⁰. Modulation of small GTPase signaling may therefore be an attractive tool to inhibit the inflammatory process in RA. In this thesis we analyzed the in vitro and in vivo effects of manipulating the activity of representative small GTPases in respect to inflammatory activation of cells relevant to RA, and animal models of this disease.

Much evidence indirectly suggests an active involvement of T cells in RA. Most importantly, the expression of the shared epitope (SE), thought to present self antigen(s) to autoreactive T cells, is one of the strongest indications of TCR involvement in RA¹¹. Synovial T cells have a highly differentiated CD45RO⁺ phenotype, promote the activation of other synovial cells through cell-cell contacts, and in many animal models T cell dependency has been shown^{12;13}. The clinical success of CTLA-4Ig therapy has suggested that TCR engagement may take place in RA, as disruption of the T cell costimulatory CD28 protein with its ligands CD80/86 is of clinical benefit¹⁴. However, despite this evidence, whether there is TCR triggering in RA is still a mat-

ter of debate. Several studies have described impaired T cell responses in RA due to deficiencies downstream of the T cell receptor (TCR). Increases in intracellular oxidative stress due to Rap1 inactivation¹⁰, impaired phosphorylation of the adaptor protein linker for activation of T cells (LAT)¹⁵ and chronic downregulation of TCRzeta expression¹⁶ are some of the defects described. Based on the inability to detect significant T cell cytokine production and proliferation in the synovium, together with the reported defects in TCR-proximal signaling responses, several studies have proposed that TCR signaling in RA is impaired¹⁷. In **chapter 2** we analyze synovial T cell responses at a single cell level, and demonstrate that TCR signaling is fully functional and completely capable of initiating T cell proliferation and cytokine production. We provide evidence that the reported hyporesponsiveness in RA synovial T cell bulk cultures is secondary to ex vivo spontaneous apoptosis. Additionally, we find that the increase in susceptibility to apoptosis ex vivo is associated with altered ratios of pro-apoptotic Noxa and anti-apoptotic Mcl-1 expression. These data indicate that although there is no evidence of T cell apoptosis in vivo, once isolated from the synovium T cells are extremely susceptible to apoptosis. It will be interesting to address which soluble factors or specific cellular contacts are responsible for the in vivo inhibition of synovial apoptosis, as abrogation of these signals might exert a therapeutic benefit by selectively promoting T cell apoptosis at the site of inflammation.

The data in **chapter 2** clearly shows that synovial T cells are capable of supporting productive TCR signaling. Our results suggest that the inability to detect T cell cytokines and proliferation in the synovium of established RA might be due to the absence of TCR engagement, rather than a consequence of TCR signaling defects. In early stages of disease, where T cell cytokines are observed, TCR engagement may play a prominent role, as autoreactive T cells could initiate inflammation. However in the effector phase of disease, T cells may contribute to inflammation by TCR-independent mechanisms. In this context it will be interesting to investigate the specific mechanism of action by which CTLA-4Ig therapy exerts its effects. In light of our results we propose that alternative mechanisms, independent of effects on classical TCR costimulation, might be responsible for the clinical improvements observed in treated patients. One of these mechanisms could be upregulation of immunomodulatory IDO on CD80/86 expressing cells. Besides antigen presenting cells (APC), T cells in the synovium have also been shown to express functional CD80/86 costimulatory molecules^{18;19}. In vitro studies have shown that triggering of CD80/86 on T cells by CTLA-4 ligation is capable of upregulating IDO expression, reducing their capacity to respond to antigen triggering²⁰. Whether T cells in the synovium, APCs, or both are targeted in this manner by CTLA-4 Ig therapy remains to be determined.

The small GTPase Rap1 is transiently activated upon TCR triggering and is an important regulator of immune responses. In RA synovial fluid (SF) T cells a block in Rap1 activation is believed to underlie the altered T cell behaviour in this disease¹⁰. While RA synovial T cells were first described as hyporesponsive, we demonstrated in **chapter 2** that T cells from the SF of RA patients are hyper-responsive to TCR triggering. In chapter 3 we investigate whether inactivation of T cell Rap1 might play an inflammatory role in RA. To accomplish this we used transgenic (Tg) mice that express constitutively active Rap1a, RapV12, within the T cell lineage. We performed a collagen-induced arthritis (CIA) experiment and evaluated the effect of Rap1 activation during experimental arthritis. We observed that arthritis incidence was decreased more than 70% in RapV12 mice and that both paw swelling as well clinical signs of disease were significantly decreased. Histological evaluation of the paws revealed that cellular infiltration and cartilage destruction were significantly reduced in Tg mice, as was joint damage. Different animal models in which the activity of Rap1 has been genetically modulated have also found defects in T cell responses. In mice expressing the active mutant Rap1E63 defects in T cell responses have been attributed to increased numbers of regulatory T cells (Tregs)²¹. In RapV12 mice we found no differences in the numbers of Tregs either under homeostatic conditions or after CIA induction, and Il-10 production was equivalent in RapV12 and WT mice. We found however, that TNF- α production was severely impaired in RapV12 mice, particularly in cytotoxic CD8⁺ T cells. Defective T cell responses have also been noted in mice lacking Spa1 expression (Spa1-/-)²². In the absence of this GAP, Rap1 activity is elevated in T cells of these mice. These mice show an age-dependent accumulation of unresponsive CD44^{high} T cells that drive defective T helper cell responses. Young Spa1^{-/-} mice with normal primary T cell responses display impaired T cell responses to recall antigens, with reduced antibody production. We investigated whether this was also the case for RapV12 mice. CD44 expression levels in RapV12 mice were comparable to WT mice and no differences were observed between naive, effector memory or central memory populations. We found however, that the circulating levels of specific anti-collagen antibody production were significantly decreased in RapV12 mice. We further investigated the mechanism underlying the defective antibody production and found that RapV12 T cells were impaired in their capacity to upregulate CD40L and ICOS co-stimulatory molecules, which could explain why RapV12 T cells are less effective in providing help for B cell activation and antibody production. Analysis of B cell responses in experimental models where T cells are dispensable for B cell activation (TNP Ficoll) and in T cell-dependent assays (TNP KLH) may be useful to confirm the defective T helper phenotype of RapV12 mice. Additionally, as CIA is a model of a chronic immune stimulation, it will be of interest to determine if the defective T cell responses in RapV12 Tg mice (lower TNF- α and antibody production) occur only in a chronic setting, or also in primary and secondary acute immune responses. Finally, it is as yet uncertain if the ability of RapV12 to protect against autoimmune disease is specific for CIA, or will be a general effect in other immune-mediated inflammatory models such as EAE.

TCR engagement in the presence of CD28 costimulation is a critical step in the initiation of effector T cell responses, while CTLA-4 ligation is essential at the end of an immune response to dampen cellular activation. Costimulation is therefore a crucial step in determining the fate of T cell responses, however the specific processes downstream of costimulatory molecule ligation are not yet clear. Accumulating studies have shown the involvement of the small GTPase Rap1 in this process. Rap1 is transiently activated upon TCR triggering and further enhanced by CTLA-4 engagement, whereas CD28 ligation blocks Rap1 activation²³⁻²⁵. These data suggests Rap1 as a central regulator of T cell responses to co-stimulatory cues. Rap1 is activated by guanine nucleotide exchange factors (GEFs) and inactivated by GTPaseactivating proteins (GAPs)²⁶. CD28-dependent inactivation of Rap1 is believed to be the result of Rap1GAP activation, while CTLA-4-dependent Rap1 activation the result of Rap1GAP inactivation²³⁻²⁵. In vivo studies have confirmed the idea of Rap1 as a critical mediator of T cell responses²⁷. Different phenotypes of transgenic (Tg) Rap1 GAP animals suggest that each Rap1 GAP may contribute differentially to T cell activation^{24;28}. In **chapter 4** we show that the five mammalian Rap1 GAPs are differentially expressed in resting and TCR/CD28 activated T cells. We found Spa-1 to be upregulated both at the mRNA and protein levels after TCR/CD28 stimulation in CD4 but not CD8 T cells. We found this upregulation to be PI3-kinase and NF-κB dependent, as inhibitors for each signaling pathway were able to block Spa-1 upregulation. We show that rather than dependent on proliferation, Spa1 upregulation is dependent on TCR/CD28 specific signals, as stimulation with IL-15, IL-17 or IL-2/ PHA were able to induce proliferation but not Spa1 expression. Additionally, we found Rap1GAP1A and Rap1GAP1B mRNA to be equally upregulated upon TCR/ CD28. Surprisingly we found the opposite at the protein level, suppression of Rap1GAP1 expression accompanied by protein degradation. Rap1GAP2 was detected at the mRNA level in rested cells, and downregulated upon TCR/CD28 stimulation. At the protein level Rap1GAP2 expression followed the same pattern as Rap1GAP1, expression in rested cells followed by suppression and protein degradation upon TCR/CD28 ligation. Distinct studies have described that post-translational regulation of Rap1GAP protein stability is necessary to avoid proteasomal degradation²⁹⁻³¹. The mechanism through which Rap1GAP proteins are degraded during T cell stimulation, and the consequences of inhibiting this process, remains to be determined.

It will be interesting to address which Rap1 GAP regulates Rap1 inactivation in CD8 T cells upon TCR/CD28, as Spa-1 upregulation occurs in CD4 T cells only. RAP-1GAP1 would be a good candidate, as it becomes upregulated at the mRNA level. Although we observed RAP1GAP1 protein degradation after 24h of TCR/CD28 stimulation, shorter time-course stimulations may elucidate on the dynamics of RAP-1GAP1 activation and degradation.

The findings presented in this chapter suggest that each Rap1 GAP may differentially contribute to T cell activation. Each Rap1 GAP may couple distinct extracellular stimuli to Rap1 regulation, associate to different costimulatory molecules and/ or inactivate Rap1 in distinct cellular subcompartments. In this chapter we study how TCR/CD28 stimulation modulates Rap1 GAP expression. It will be interesting to evaluate how other co-stimulatory molecules, either positive (ICOS) or negative (CTLA-4 and PD-1) regulators of T cell activation influence Rap1 activation. Investigating the activation patterns of each Rap1GAP may allow to the development of compounds that will be able to specifically regulate Rap1 activation in distinct T cell compartments, with potential application to diseases like RA where a constitutive block in Rap1 expression is observed.

Fibroblast-like synoviocytes (FLS) from RA patients display proliferative and invasive properties reminiscent of malignant tumor cells³². Ras small GTPases play important roles in tumor cell proliferation and invasion. Although no transforming mutations have been detected in RA FLS^{33,34}, pathways mediated by Ras GTPases are known to be de-regulated^{7;35;36}. Recent evidence suggests that changes in GEF expression or activity might be relevant to disease. This is the case in human melanoma cells, where expression levels of Ras guanine nucleotide-releasing factor (RasGRF) 1 regulates constitutive MMP-9 production³⁷. In **chapter 5** we analyze the expression patterns of RasGRF1, a specific H-Ras activator³⁸, in RA and non-RA ST and FLS. RasGRF1 expression was significantly increased in RA ST and found to positively correlate and co-localize with MMP-1 and MMP-3 producing cells. Additionally, we found that in vitro modulation of RasGRF1 expression in RA FLS was able to regulate spontaneous MMP-3 production. We could not find an association between RasGRF1 expression and MMP-1 production in RA FLS, which could imply that maybe other cell types, such as macrophages might be more important sources of MMP-1 production. It will therefore be interesting to analyze the effects of knock down and overexpression of RasGRF1 in macrophages in terms of MMP-1 production. In lysates from FLS and ST we observed the expression of post-translationally modified RasGRF1. These truncated fragments are known to be derived from protease-dependent cleavage and these fragments have been shown to have enhanced Ras-activating capacity³⁹. The results presented in this chapter suggest that enhanced RasGRF1 expression and activating post-translational modifications in RA FLS sensitize signaling pathways that promote MMP-3 production and consecutively joint damage. It is not yet known which stimuli present in the joint promotes the post-translational modification of RasGRF1, but approaches targeting these activating stimuli may prove of clinical interest in reducing RasGRF1 activation and MMP-3 production.

Cellular migration, proliferation, activation and immunological synapse formation are processes tightly regulated by Rho family Rac small GTPases⁴⁰. In RA these cellular processes are thought to be de-regulated as high numbers of activated immune cells are observed in the synovium, reflecting increased migration into the joints, local proliferation or both⁴¹. In **chapter 6** we investigated whether treatment of arthritic mice with inhibitory Rac1 C-terminal peptide, which specifically blocks Rac1 signaling, would ameliorate disease severity. Different studies have reported the involvement of members of the Rho GTPase family in the pathogenesis of RA. Inhibition of Rac1 signaling in vitro is able to reduce FLS proliferation and invasiveness⁸ while modulation of Rho activation regulates FLS proliferation and IL-6 secretion⁹. In vivo, in a model of reactive arthritis, Rac signaling in neutrophils has been shown to regulate joint inflammation⁴². In this chapter we treated arthritic mice during early or chronic disease, with Rac1 C-terminal peptide and found that inhibition of Rac1 signaling in vivo significantly reduces anti-collagen antibody production. The reduction in levels of immunoglobulin (Ig) G1 and IgG2a (early treatment) or IgG2a and IgG2b (treatment during chronic arthritis) correlated with an amelioration of paw swelling in treated animals. Decreased antibody production could be the result of defective T cell priming, as B cell responses are critically dependent on proper T cell activation. In order to provide help for B cell activation, T cells need to have a productive engagement with APC. Studies in vivo have shown that Rac deficient dendritic cells (DC) have defective migration to secondary lymphoid organs and fail to establish stable contacts with naive T cells⁴³. In this case, suboptimal T cell priming leads to defective humoral responses to T cell-dependent antigens⁴⁴. Rac1 peptide may similarly inhibit anti-collagen antibody production by disturbing T cell-APC interactions. Further studies addressing antibody production in T cell -dependent and -independent assays may elucidate whether treatment with Rac1 C-terminal peptide targets T cell priming or the antibody producing B cells directly. In the first case, it will be interesting to address whether also T cell effector responses, besides T helper, are impaired.

The observed effect on the reduction of paw swelling could be due to effects in edema formation, as inhibition of Rac1-mediated signaling leads to a decrease in leukocyte extravasation⁴⁵. However, although in vitro Rac1 C-terminal peptide was able to inhibit actin polymerization, we found no evidence for a decrease in cellular infiltration or joint destruction in vivo. The limited clinical effect of Rac1 peptide treatment may reflect the redundant role of Rac2 in promoting osteoclastogenene-sis^{46,47} and cellular migration⁴⁸, as the inhibitory peptide we used is highly specific for Rac1 and does not interfere with Rac2 signaling⁴⁹. Although the data presented in this chapter shows that treatment of arthritic mice with Rac1 peptide can safely be used to reduce paw swelling and the circulating levels of anti-collagen antibodies, future studies will need to directly evaluate the relative contributions of both Rac1 and Rac2 in CIA.

In conclusion, the work presented in this thesis shows that modulation of small GTPase signaling might have good therapeutic potential in the treatment of RA. Strategies aimed at the inhibition or maintenance of small GTPase signaling could be applied at different levels of intracellular signaling cascades and tested for their efficacy in the amelioration of arthritis.

Several compounds targeting effectors of small GTPases are already available, and some have been tested in clinical trials with successful results. In cancer clinical trials, blockage of the Raf/MEK/ERK pathway with compounds targeting Raf (BAY 43-9006, ISIS 5132) and MEK (PD184352, PD0325901 and ARRY-142886) have showed promising results⁵⁰. In RA clinical trials, the use of VX-702 and RO-440-2257 as p38 inhibitors have showed only limited success^{51,52}, and further analysis of the distinct contributions of Ras effector pathways to RA is urgently needed.

Direct targeting of small GTPases has also been investigated in different disease settings. We have shown in **chapter 6** the possibility of using C-terminal peptides to interfere with specific small GTPase signaling in vivo for the treatment of CIA. In chapter 5 we used locked nucleic acid (LNA) as a tool to efficiently inhibit RasGRF1 expression in vitro. Compounds that interfere with small GTPase post-translational modification, such as farnesyl transferase inhibitors (FTIs), have received special attention in cancer settings. The FTIs Tipifarnib and lonafarnib are currently under clinical investigation with encouraging results in the treatment of hematological malignancies and breast cancer⁵³. Although not yet tested in clinical trials, bacterial toxins have also been shown to target individual small GTPases. Clostridium difficile toxin A and B specifically glucosylate Rho proteins, while bacterial C3-like ADP-ribosyltransferase induces ADP-ribosylation of RhoA, both inhibiting in a specific manner small GTPase downstream signaling^{54,55}. Maintenance or activation of small GTPases may also be interesting to modulate inflammation in RA. We have shown in **chapter 3** that maintenance of Rap1 signaling ameliorates disease in CIA. 8-(4-chloro-phenylthio)-2'-O-methyladenosine-3',5'-cyclic monophosphate (8CPT-2Me-cAMP) is a cAMP analog that can selectively activate Epac, a Rap1 GEF⁵⁶. Although not yet tested in a disease setting, it would be interesting to analyze the effect of this analogue in mice with CIA.

Finally, targeting cell surface receptors upstream of the small GTPases may allow specific modulation of GTPase activity, through the regulation of GEFs and GAPs. One such candidate may already be in the clinic. CTLA-4Ig (abatacept), which disrupts the interaction between the co-stimulatory CD28 molecule and CD80/86 on APCs, has shown clinical benefit in RA patients. In vitro, this compound has been shown to prevent T cell Rap1 inactivation by synovial fluid adherent cells⁵⁷. Future work will need to determine how restored T cell Rap1 function may contribute to the clinical benefit of CTLA-4Ig therapy. In sum, the data presented in this thesis provides strong pre-clinical evidence that targeting Ras superfamily GTPases may be beneficial in the treatment of RA, a potential awaiting the development of compounds specifically targeting these enzymes in the clinic.

References

- 1. Olsen NJ, Stein CM. New drugs for rheumatoid arthritis. N.Engl.J.Med. 2004;350:2167-2179.
- Schett G, Tohidast-Akrad M, Smolen JS et al. Activation, differential localization, and regulation of the stress-activated protein kinases, extracellular signalregulated kinase, c-JUN N-terminal kinase, and p38 mitogen-activated protein kinase, in synovial tissue and cells in rheumatoid arthritis. Arthritis Rheum. 2000;43:2501-2512.
- 3. Tak PP, Firestein GS. NF-kappaB: a key role in inflammatory diseases. J.Clin. Invest 2001;107:7-11.
- 4. Zhang HG, Wang Y, Xie JF et al. Regulation of tumor necrosis factor alpha-mediated apoptosis of rheumatoid arthritis synovial fibroblasts by the protein kinase Akt. Arthritis Rheum. 2001;44:1555-1567.
- Na HJ, Lee SJ, Kang YC et al. Inhibition of farnesyltransferase prevents collagen-induced arthritis by down-regulation of inflammatory gene expression through suppression of p21(ras)-dependent NF-kappaB activation. J.Immunol 2004;173:1276-1283.
- 6. Pap T, Nawrath M, Heinrich J et al. Cooperation of Ras- and c-Myc-dependent pathways in regulating the growth and invasiveness of synovial fibroblasts in rheumatoid arthritis. Arthritis Rheum. 2004;50:2794-2802.
- 7. Yamamoto A, Fukuda A, Seto H et al. Suppression of arthritic bone destruction by adenovirus-mediated dominant-negative Ras gene transfer to synoviocytes and osteoclasts. Arthritis Rheum. 2003;48:2682-2692.
- 8. Chan A, Akhtar M, Brenner M et al. The GTPase Rac regulates the proliferation and invasion of fibroblast-like synoviocytes from rheumatoid arthritis patients. Mol.Med. 2007;13:297-304.
- 9. Nakayamada S, Kurose H, Saito K, Mogami A, Tanaka Y. Small GTP-binding protein Rho-mediated signaling promotes proliferation of rheumatoid synovial fibroblasts. Arthritis Res.Ther. 2005;7:R476-R484.
- 10. Remans PH, Gringhuis SI, van Laar JM et al. Rap1 signaling is required for suppression of Ras-generated reactive oxygen species and protection against oxidative stress in T lymphocytes. J.Immunol. 2004;173:920-931.
- 11. Winchester R, Dwyer E, Rose S. The genetic basis of rheumatoid arthritis. The shared epitope hypothesis. Rheum.Dis.Clin.North Am. 1992;18:761-783.
- 12. Lundy SK, Sarkar S, Tesmer LA, Fox DA. Cells of the synovium in rheumatoid

arthritis. T lymphocytes. Arthritis Res. Ther. 2007;9:202.

- 13. Thomas R, McIlraith M, Davis LS, Lipsky PE. Rheumatoid synovium is enriched in CD45RBdim mature memory T cells that are potent helpers for B cell differentiation. Arthritis Rheum. 1992;35:1455-1465.
- 14. Genovese MC, Schiff M, Luggen M et al. Efficacy and safety of the selective costimulation modulator abatacept following 2 years of treatment in patients with rheumatoid arthritis and an inadequate response to anti-tumour necrosis factor therapy. Ann.Rheum.Dis. 2008;67:547-554.
- 15. Gringhuis SI, Leow A, Papendrecht-van der Voort EA et al. Displacement of linker for activation of T cells from the plasma membrane due to redox balance alterations results in hyporesponsiveness of synovial fluid T lymphocytes in rheumatoid arthritis. J.Immunol. 2000;164:2170-2179.
- 16. Zhang Z, Gorman CL, Vermi AC et al. TCRzetadim lymphocytes define populations of circulating effector cells that migrate to inflamed tissues. Blood 2007;109:4328-4335.
- 17. Firestein GS, Zvaifler NJ. How important are T cells in chronic rheumatoid synovitis?: II. T cell-independent mechanisms from beginning to end. Arthritis Rheum. 2002;46:298-308.
- Azuma M, Yssel H, Phillips JH, Spits H, Lanier LL. Functional expression of B7/ BB1 on activated T lymphocytes. J.Exp.Med. 1993;177:845-850.
- 19. Verwilghen J, Lovis R, De BM et al. Expression of functional B7 and CTLA4 on rheumatoid synovial T cells. J.Immunol. 1994;153:1378-1385.
- 20. Boasso A, Herbeuval JP, Hardy AW, Winkler C, Shearer GM. Regulation of indoleamine 2,3-dioxygenase and tryptophanyl-tRNA-synthetase by CTLA-4-Fc in human CD4+ T cells. Blood 2005;105:1574-1581.
- 21. Li L, Greenwald RJ, Lafuente EM et al. Rap1-GTP is a negative regulator of Th cell function and promotes the generation of CD4+CD103+ regulatory T cells in vivo. J.Immunol. 2005;175:3133-3139.
- 22. Ishida D, Yang H, Masuda K et al. Antigen-driven T cell anergy and defective memory T cell response via deregulated Rap1 activation in SPA-1-deficient mice. Proc.Natl.Acad.Sci.U.S.A 2003;100:10919-10924.
- 23. Carey KD, Dillon TJ, Schmitt JM et al. CD28 and the tyrosine kinase lck stimulate mitogen-activated protein kinase activity in T cells via inhibition of the small G protein Rap1. Mol.Cell Biol. 2000;20:8409-8419.
- 24. Dillon TJ, Carey KD, Wetzel SA, Parker DC, Stork PJ. Regulation of the small GTPase Rap1 and extracellular signal-regulated kinases by the costimulatory molecule CTLA-4. Mol.Cell Biol. 2005;25:4117-4128.

- 25. Reedquist KA, Bos JL. Costimulation through CD28 suppresses T cell receptordependent activation of the Ras-like small GTPase Rap1 in human T lymphocytes. J.Biol Chem. 1998;273:4944-4949.
- 26. Bos JL, Rehmann H, Wittinghofer A. GEFs and GAPs: critical elements in the control of small G proteins. Cell 2007;129:865-877.
- 27. Minato N, Kometani K, Hattori M. Regulation of immune responses and hematopoiesis by the Rap1 signal. Adv.Immunol. 2007;93:229-264.
- 28. Kometani K, Moriyama M, Nakashima Y et al. Essential role of Rap signal in pre-TCR-mediated beta-selection checkpoint in alphabeta T-cell development. Blood 2008;112:4565-4573.
- 29. Gao Q, Srinivasan S, Boyer SN, Wazer DE, Band V. The E6 oncoproteins of highrisk papillomaviruses bind to a novel putative GAP protein, E6TP1, and target it for degradation. Mol.Cell Biol. 1999;19:733-744.
- 30. Jordan JD, He JC, Eungdamrong NJ et al. Cannabinoid receptor-induced neurite outgrowth is mediated by Rap1 activation through G(alpha)o/i-triggered prote-asomal degradation of Rap1GAPII. J.Biol.Chem. 2005;280:11413-11421.
- 31. Tsygankova OM, Feshchenko E, Klein PS, Meinkoth JL. Thyroid-stimulating hormone/cAMP and glycogen synthase kinase 3beta elicit opposing effects on Rap1GAP stability. J.Biol.Chem. 2004;279:5501-5507.
- 32. Tak PP, Zvaifler NJ, Green DR, Firestein GS. Rheumatoid arthritis and p53: how oxidative stress might alter the course of inflammatory diseases. Immunol Today 2000;21:78-82.
- 33. Pap T, Franz JK, Hummel KM et al. Activation of synovial fibroblasts in rheumatoid arthritis: lack of Expression of the tumour suppressor PTEN at sites of invasive growth and destruction. Arthritis Res. 2000;2:59-64.
- 34. Roivainen A, Zhu F, Sipola E, Yli-Jama T, Toivanen P. Failure to verify H-ras mutations in arthritic synovium: comment on the article by Roivainen et al. Arthritis Rheum. 2001;44:2705.
- 35. Na HJ, Lee SJ, Kang YC et al. Inhibition of farnesyltransferase prevents collagen-induced arthritis by down-regulation of inflammatory gene expression through suppression of p21(ras)-dependent NF-kappaB activation. J.Immunol. 2004;173:1276-1283.
- 36. Pap T, Nawrath M, Heinrich J et al. Cooperation of Ras- and c-Myc-dependent pathways in regulating the growth and invasiveness of synovial fibroblasts in rheumatoid arthritis. Arthritis Rheum. 2004;50:2794-2802.
- 37. Zhu TN, He HJ, Kole S et al. Filamin A-mediated down-regulation of the exchange factor Ras-GRF1 correlates with decreased matrix metalloproteinase-9

expression in human melanoma cells. J.Biol.Chem. 2007;282:14816-14826.

- Jones MK, Jackson JH. Ras-GRF activates Ha-Ras, but not N-Ras or K-Ras 4B, protein in vivo. J.Biol.Chem. 1998;273:1782-1787.
- Baouz S, Jacquet E, Bernardi A, Parmeggiani A. The N-terminal moiety of CDC25(Mm), a GDP/GTP exchange factor of Ras proteins, controls the activity of the catalytic domain. Modulation by calmodulin and calpain. J.Biol.Chem. 1997;272:6671-6676.
- 40. Jaffe AB, Hall A. Rho GTPases: biochemistry and biology. Annu.Rev.Cell Dev. Biol. 2005;21:247-269.
- 41. Buckley CD. Michael Mason prize essay 2003. Why do leucocytes accumulate within chronically inflamed joints? Rheumatology.(Oxford) 2003;42:1433-1444.
- 42. Zhang X, Glogauer M, Zhu F et al. Innate immunity and arthritis: neutrophil Rac and toll-like receptor 4 expression define outcomes in infection-triggered arthritis. Arthritis Rheum. 2005;52:1297-1304.
- 43. Benvenuti F, Hugues S, Walmsley M et al. Requirement of Rac1 and Rac2 expression by mature dendritic cells for T cell priming. Science 2004;305:1150-1153.
- 44. Li H, Oliver T, Jia W, He YW. Efficient dendritic cell priming of T lymphocytes depends on the extracellular matrix protein mindin. EMBO J. 2006;25:4097-4107.
- 45. van Wetering S, van den BN, van Buul JD et al. VCAM-1-mediated Rac signaling controls endothelial cell-cell contacts and leukocyte transmigration. Am.J.Physiol Cell Physiol 2003;285:C343-C352.
- Fukuda A, Hikita A, Wakeyama H et al. Regulation of osteoclast apoptosis and motility by small GTPase binding protein Rac1. J.Bone Miner.Res. 2005;20:2245-2253.
- 47. Wang Y, Lebowitz D, Sun C et al. Identifying the relative contributions of Rac1 and Rac2 to osteoclastogenesis. J.Bone Miner.Res. 2008;23:260-270.
- 48. Sun CX, Downey GP, Zhu F et al. Rac1 is the small GTPase responsible for regulating the neutrophil chemotaxis compass. Blood 2004;104:3758-3765.
- 49. Van Hennik PB, ten Klooster JP, Halstead JR et al. The C-terminal domain of Rac1 contains two motifs that control targeting and signaling specificity. J.Biol. Chem. 2003;278:39166-39175.
- 50. Kohno M, Pouyssegur J. Targeting the ERK signaling pathway in cancer therapy. Ann.Med. 2006;38:200-211.
- 51. Damjanov N, Kauffman RS, Spencer-Green GT. Efficacy, pharmacodynamics, and safety of VX-702, a novel p38 MAPK inhibitor, in rheumatoid arthritis: results of two randomized, double-blind, placebo-controlled clinical studies.

Arthritis Rheum. 2009;60:1232-1241.

- 52. Cohen SB, Cheng TT, Chindalore V et al. Evaluation of the efficacy and safety of pamapimod, a p38 MAP kinase inhibitor, in a double-blind, methotrexate-controlled study of patients with active rheumatoid arthritis. Arthritis Rheum. 2009;60:335-344.
- 53. Karp JE, Lancet JE. Development of farnesyltransferase inhibitors for clinical cancer therapy: focus on hematologic malignancies. Cancer Invest 2007;25:484-494.
- 54. Just I, Selzer J, Wilm M et al. Glucosylation of Rho proteins by Clostridium difficile toxin B. Nature 1995;375:500-503.
- 55. Fritz G, Kaina B. Rho GTPases: promising cellular targets for novel anticancer drugs. Curr.Cancer Drug Targets. 2006;6:1-14.
- 56. Enserink JM, Christensen AE, de RJ et al. A novel Epac-specific cAMP analogue demonstrates independent regulation of Rap1 and ERK. Nat.Cell Biol. 2002;4:901-906.
- 57. Remans PH, Wijbrandts CA, Sanders ME et al. CTLA-4IG suppresses reactive oxygen species by preventing synovial adherent cell-induced inactivation of Rap1, a Ras family GTPASE mediator of oxidative stress in rheumatoid arthritis T cells. Arthritis Rheum. 2006;54:3135-3143.