*Downloaded from UvA-DARE, the institutional repository of the University of Amsterdam (UvA)*
*http://dare.uva.nl/document/161387*

| | |
|---|---|
| File ID | 161387 |
| Filename | References |

SOURCE (OR PART OF THE FOLLOWING SOURCE):

| | |
|---|---|
| Type | Dissertation |
| Title | Symbiosis of thrips and gut bacteria |
| Author | E.J. de Vries |
| Faculty | Faculty of Science |
| Year | 2010 |
| Pages | 173 |
| ISBN | 978 90 76894 85 0 |

FULL BIBLIOGRAPHIC DETAILS:
http://dare.uva.nl/record/328135

*Copyright*

*It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use.*

*UvA-DARE is a service provided by the library of the University of Amsterdam (http://dare.uva.nl)*

R

# References


Adams, D., and Douglas, A. E. (1997) How symbiotic bacteria influence plant utilisation by the polyphagous aphid, *Aphis fabae*. Oecologia 110:528-532.

Akman, L., Yamashita, A., Watanabe, H., Oshima, K., Shiba, T., Hattori, M., and Aksoy, S. (2002) Genome sequence of the endocellular obliate symbiont of tsetse flies, *Wigglesworthia glossinida.* Nature Genetics 32:402-407.

Aksoy, S., Pourhoseini, A., and Chow, A. (1995) Mycetome endosymbionts of tsetse flies constitute a distinct lineage related to Enterobacteriaceae. Insect Mol. Biol. 4:15-22.

Ammar, E. D, (1994) Propagative transmission of plant and animal viruses by insects: Factors affecting vector specificity and competence. Adv. Dis. Vector Res. 10:289-331.

Andrews, R. E., and Spence, K. D. (1980) Action of Douglas Fir Tussock Moth larvae and their microflora on dietary terpenes. Appl. Environ. Microbiol. 40:959-963.

Baerwald, R. J., and Boush, G. M. (1968) Demonstration of the bacterial symbiote *Pseudomonas mellophtera* in the Apple Maggot, *Rhagoletis pomonella*, by the fluorescent-antibody technique. J. Invertebr. Pathol. 11:251-259.

Baines, S.(1956) The role of the symbiotic bacteria in the nutrition of *Rhodnius prolixus* (Hemiptera). J. Exp. Biol 33:533-541.

Bandi, C., Damiani, G., Magrassi, L., Grigilo, A., Fani, R., and Sacchi, L. (1994) Flavobacteria as intracellular symbionts in cockroaches. Proc. R. Soc. London B. 257:43-48.

Bandi, C., Sironi, M., Damiani, G., Magrassi, L., Nalepa, C. A., Laudani, U., and Sacchi, L. (1995) The establishment of intracellular symbiosis in an ancestor of cockroaches and termites. Proc. R. Soc. London B 259:293-299.

Bauchop, T., and Clarke, R. T. J. (1975) Gut microbiology and carbohydrate digestion in the larva of *Costelytra zeelandica* (Coleoptera: Scarabeidae). New Zeal. J. Zool. 2:237-243.

Baumann, L., and Baumann, P. (1994) Growth kinetics of the endosymbiont *Buchnera aphidicola* in the aphid *Schizaphis graminum*. Appl. Environ. Microb. 60:3440-3443.

Baumann, P., Baumann, L., Lai, C.-Y., Rouhbakhsh, D., Moran, N. A., and Clark, M. A. (1995) Genetics, physiology, and evolutionary relationships of the genus *Buchnera*, intracellular symbionts of aphids. Annu. Rev. Microbiol. 49:55-94.

Baumann, P., and Moran, N. A. (1997) Non-cultivable microorganisms from symbiotic associations of insects and other hosts. Anth. Leeuwenh. 72:39-48.

Bayon, C. (1980) Volatile fatty acids and methane production in relation to anaerobic carbohydrate fermentation in *Oryctes nasicornis* larvae (Coleoptera: Scarabeidae). J. Insect Physiol. 26:819-828.

Beard, C. B., Cordon-Rosales, C., and Durvasula, R. V. (2002) Bacterial symbionts of the Triatomine and their potential use in control of Chagas disease transmission. Annu. Rev. Entomol. 47:123-141.

Behar, A., Yuval, B., and Jurkevitz, E. (2005) Enterobacteria-mediated nitrogen fixation in natural pop-

ulations of the fruit fly Ceratitis capitata. Mol. Ecol. 14:2637-2643.

Behrenz, W., and Technau, G. (1959) Versuche zur Bekampfung von *Anobium bipunctatum* mit Symbionticiden. Z. Angew. Entomol. 4:22-28.

Beijerinck, M. W., and Rant, A. (1905) Wundreiz, Parasitismus und Gummifluss bei den Amygdaleen. Zentralbl. Bakteriol. Parasistenk. Infektionskr. Abt II 11:366-375.

Belliure, B., Janssen, A., Maris, P. C., Peters, D., and Sabelis, M. W. (2005) Herbivore arthropods benefit from vectoring plant viruses. *Ecol. Let.* 8:70-79.

Berenbaum, M. R. (1988) Micro-organisms as mediators of intertrophic and intratrophic interactions. *In:* Novel Aspects of Insect-Plant Interactions (P. Barbosa and D. K. Letourneau eds.), pp. 91-123. Plenum press, New York (N.Y.), USA.

Bergey, D. H. (ed.) (1974) the Bergey's Manual of Determinative Bacteriology. .

Bernays, E. A., and Graham, M. (1988) On the evolution of host specificity in phytophagous arthropods. Ecol. 69:886-892.

Bewig, F., and Schwartz, W. (1956) Untersuchungen über die Symbiose von Tieren mit Pilzen und Bakterien. VII Über die Physiologie der Symbiose bei einigen blutsauchenden Insekten. Arch. Microbiol. 24:174-208.

Bignell, D. E. (1983) The arthropod gut as an environment for microorganisms. *In*: Microbial Ecology of the Gut (R. T. J. Clarke and T. Bauchop, eds.), pp. 205-227. Acad. Press, New York, NY, USA.

Bignell, D. E., Oskarsson, H., and Anderson, J. M. (1980) Distribution and abundance of bacteria in the gut of a soil-feeding termite *Procurbitermes aburiensis* (Termitidae, Termitinae). J. Gen. Microbiol. 117:393-403.

Boucher, D. H., James, S., and Keeler, K. H. (1982) The ecology of mutualism. Annu. Rev. Ecol. Syst. 13:315-347.

Bournier, A. (1961) Sur l'existence et l'évolution d'un mycétome au cours de l'embryogenèse de *Caudothrips buffai* Karny. *In:* Verh. XI Int. Kong. Entomol, pp. 352-354.

Boush, G. M., Saleh, S. M., and Baranovski, R. M. (1972) Bacteria associated with the Caribbean fruit fly. Environ. Entomol. 1:30-33.

Boush, G. M., and Matsumura, F. (1967) Insecticidal degradation by *Pseudomonas melophtora*, the bacterial symbiote of the apple maggot. J. Econ. Entomol. 60:918-920.

Bracke, J. W., Cruden, D. L., and Markovetz, A. J. (1979) Intestinal microbial flora of the American cockroach, *Periplaneta americana* L. Appl. Environ. Microbiol. 38:945-955. .

Brand, J. M., Bracke, J. W., Markovetz, A. J., Wood, D. L., and Browne, L. E. (1975) Production of verbenol pheromone by a bacterium isolated from bark beetles. Nature 254:136-137.

Brauman, A., Doré, J., Eggleton, P., Bignell, D., Breznak, J. A., and Kane, M. D. (2001) Molecular phylogenetic profiling of prokaryotic communities in guts of termites with different feeding habits. FEMS Microb. Ecol. 35:27-36.

Breeuwer, J. A. J., Stouthamer, R., Barnes, S. M., Pelletier, D. A., Weisburg, W. G., and Werren, J. H. (1992) Phylogeny of cytoplasmic incompatibility microorganisms in the parasitoid wasp genus *Nasonia*. Insect Mol. Biol. 1:25-36. .

Breeuwer, J. A. J., and Werren, J. H. (1993) Cytoplasmic incompatibility and bacterial density in *Nasonia vitripennis*. Genetics 135: 565-574.

Breeuwer, J. A. J. (1997) *Wolbachia* and cytoplasmic incompatibility in the spider mites *Tetranychus urticae* and *T. turkestani*. Heredity 79: 41-47.

Breznak, J. A. (1982) Intestinal microbiota of termites and other xylophagous insects. Annu. Rev. Microbiol. 36:323-343.

Breznak, J. A., and Brune, A. (1994) Role of microorganisms in the digestion of lignocellulose by ter-

mites. Annu. Rev. Entomol. 39:453-487.

Bridges, J. R. (1981) Nitrogen-fixing bacteria associated with bark beetles. Microbial Ecol. 7:131-137.

Broderick, N. A., Raffa, K. F., Goodman, R. M., and Handelsman, J. (2004) Census of the bacterial community of the gypsy moth larval midgut by using culturing and culture-independent methods. Appl. Environ. Microb. 70:293-300.

Brødsgaard, H. F. (1989) *Frankliniella occidentalis* (Thysanoptera: Thripidae) – a new pest in Danish glasshouses. A review. Tidsskrift Planteavl. 93:83-91.

Brødsgaard, H. F. (1994) Insecticide resistance in European and African strains of western flower thrips (Thysanoptera: Thripidae) tested in a new residue-on-glass test. J. Econ. Entomol. 87:1141-1146.

Bronstein, J. L. (1994a) Conditional outcomes of mutualistic interactions, Trends Ecol. Evol. 9: 214-217.

Bronstein, J. L. (1994b) Our current understanding of mutualism. Quat. Rev. Biol. 69:31-51.

Brune, A., Emerson, D., and Breznak, J. A. (1995) The termite gut microflora as an oxygen sink: microelectrode determination of oxygen and pH gradients in guts of lower and higher termites. Appl. Environ. Microbiol. 61:2681-2687.

Brunner, P. C., and Frey, J. E. (2004) Isolation and characterization of six polymorphic loci in the western flower thrips *Frankliniella occidentalis* (Insecta, Thysanoptera). Molec. Ecol. Notes 4:599-600.

Brunner, P. C., Chatzivassiliou, E. K., Katis, N. I. , and Frey, J. E. (2004) Host-associated genetic differentiation in *Thrips tabaci* (Insecta; Thysanoptera), as determined from mtDNA sequence data. Heredity 93:364-370.

Bryan, D. E., and Smith, R. F. (1956) The *Frankliniella occidentalis* (Pergande) complex in California (Thysanoptera: Thripida), p. 350-410. University of California press Berkeley and Los Angeles, CA, USA.

Bucher, G. E., and Stephens, J. M. (1959) Bacteria of grasshoppers. J. Invertebr. Pathol. 1:365-390. .

Buchner, P. (1920) Zur Kenntnis der Symbiose niederer pflanzlicher Organismen mit Pedikuliden, Biol. Zentralbl. 39:109-112.

Buchner, P. (1953) “Symbiose der Tiere mit pflanzliche Mikroorganismen”. Birkhäuser, Basel (Switzerland).

Buchner, P. (1965) Endosymbiosis of Animals with Plant Microorganims. John Wiley and sons, New York, NY, USA. 905 pp.

Burges, H. D., Grove, J. F., and Pople, M. (1979) The internal microbial flora of the elm bark beetle, *Scolytus scolytus*, at all stages of its development. J. Invertebr. Pathol. 34:21-25. .

Bush, G. L., and Smith, J. J. (1998) Genetics and ecology of sympatric speciation, a case study. Res. in Popul. Ecol. 40:175-187.

Campbell, B. C. (1990) On the role of microbial symbiotes in herbivorous insects. *In:* Insect-Plant Interactions (E. A. Bernays, ed.), pp. 1-45. CRC press, Boca Raton, LA, USA.

Campbell, B. C., Bragg, T. S., and Turner, Ch. (1992) Phylogeny of symbiotic bacteria of four weevil species (Coleoptera: Curculionidae) based on analysis of 16S ribosomal DNA. Insect Biochem. Mol. Biol. 22:415-421.

Cazemier, A. E., Hackstein, J. H. P., Op den Camp, H. J. M., and Van der Drift, Chr. (1997) Bacteria in the instinal tract of different species of arthropods. Microb. Ecol. 33:189-197.

Cazemier, A. E., Verdoes, J. C., Ruebsat, F. A. G., Hackstein, J. H. P., Van der Drift, Chr., and Op den Camp, H. J. M. (2003) *Promicromonospora pachnodae* sp. nov., a member fo the (hemi)cellulolytic hindgut flora of larvae of the scarab beetle *Pachnoda marginata*. Anth. Van Leeuwenhoek 83:135-148.

Chang, K. P. (1974) Effects of elevated temperature on the mycetome and symbionts of the bed bug *Cimex lectularius* (Heteroptera). J. Invertebr. Pathol. 23:333-340.

Chang, K. P., and Musgrave, A. J. (1972) Multiple symbiosis in a leafhopper, *Helochara communis* Fitch (Cicadellidae: Homoptera): envelopes, nucleoids and inclusions of the symbiotes. J. Cell Sci. 11:275-293.

Charpentier, R., Charpentier, B., and Zehtner, O. (1978) The bacterial flora of the midgut of two Danish populations of healthy fifth instar larvae of the turnip moth, *Scotia segetum*. J. Invertebr. Pathol. 32:59-63.

Chatzivassiliou, E. K., Nagata, T., Katis, N. J., and Peters, D. (1999) Transmission of tomato spotted wilt virus by *Thrips tabaci* populations originating from leek. Plant Pathol. 48:700-706.

Chen, D.-Q., and Purcell, A. (1997) Occurrence and transmission of facultative endosymbionts in aphids. Curr. Microbiol. 34:220-225.

Chen, X., Changben, L., Zhao, S., and Aksoy, S. (1999) Phylogeny of the symbionts of tsetse and the evolutionary relationships with their hosts. Progress in Natural Sci. 9:922-928.

Clark, M. A. Baumann, L., Munson, M. A., Baumann, P., Campbell, B. C., Duffus, J. E., Osborne, L. S., and Moran, N. A. (1992) The eubacterial endosymbionts of whiteflies (Homoptera, Aleyrodoidea) constitute a lineage distinct from the endosymbionts of aphids and mealybugs. Curr. Microbiol. 25:119-123.

Claridge, M. F., and Den Hollander, J. (1983) The biotype concept and its application to insect pests of agriculture. Crop protection 2:85-95.

Cochran, D. G. (1985) Nitrogen excretion in cockroaches. Annu. Rev. Entomol. 30:29-49.

Cole, R. A., Rollason, S. A., Hardman, J. A., and Ellis, P. R. (1990) The incidence of seed-borne bacteria in carrots and their association with the damage caused by the larvae of the carrot fly (*Psila rosae*). Brighton crop protection conference - pests and diseases, pp. 343-347.

Colgan, D. J. (1987) Host-parasite genome relationships in Acridid grasshoppers. I. Variation in the isoenzymes of the *Caledia captiva* species complex and in the gut bacterium *Enterobacter cloaca*. Genome 29:257-263.

Coudriet, P. L., Kisabo, A. N., McCright, J. D., and Bohn, G. W. (1979) Varietal resistance to onion thrips. J. Econ. Entomol. 72:614-615.

Crespi, B., Carmean, D., Vawter, L., and Von Dohlen, C. (1996) Molecular phylogenetics of Thysanoptera. Syst. Entomol. 21:79-87.

Cruden, D. L., and Markovetz, A. J. (1984) Microbial aspects of the cockroach hindgut. Arch. Microbiol. 138:131-139.

Czolij, R., Slaytor, M., and O'Brien, R. W. (1985) Bacterial flora of the mixed segment and the hindgut of the higher termite *Nasutitermes exositus* Hill (Termitidae, Nasutitermitinae). Appl. Environ. Microbiol. 49:1226-1236.

Dadd, R. H. (1975) Alkalinity within the midgut of mosquito larvae with alkaline-active digestive enzymes. J. Invertebr. Pathol. 21:1847-1853.

Dallai, R., Del Bene, G., and Marchini, D. (1991) The ultrastructure of Malpighian tubules and hindgut of *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae). Int. J. Insect Morphol. Embryol. 20:223-233.

Dale, C., and Maudlin, I. (1999) *Sodalis* gen. nov. and *Sodalis glissinidis* sp. nov., a microaerophilic secondary symbiont of the tsetse fly *Glossina morsitans morsitans*. Int. J. Syst. Bacteriol. 49:267-275.

Darby, A. C., and Douglas, A. E. (2003) Elucidation of the transmission patterns of an insect-borne bacterium. Appl. Environ. Microbiol. 69:4403-4407.

Daser, U., and Brandl, R. (1992) Microbial gut floras of eight species of tephritids. Biol. J. Linn. Soc. 45:155-165.

De Ávila, A. C., Huguenot, C., Resende, R. O., Kitajima, E. W., Goldbach, R. W., and Peters, D. (1990) Serological differentiation of 20 isolates of tomato spotted wilt virus. J. Gen. Virol. 71:2801-2807.

De Bary, A. (1879) Die Erscheinung der Symbiose. Naturforsch. Versamml. Cassel, LI, Tageblatt, 121 pp.

De Jager, K. C. M. (1995) Mechanisms of Resistance to Western Flower Thrips in Chrysanthemum. Ph D thesis. Leiden University. 132 pp.

De Kogel, W. J. (1997) Host Plant Resistance to Western Flower Thrips: Variable Plants and Insects. Ph D thesis, Univ. of Amsterdam. 126 pp.

De Kogel, W. J., Van der Hoek, M., and Mollema, C. (1997a) Variation in performance of western flower thrips on susceptible and partially resistant cucumber. Entomol. Exp. Appl. 83:73-80.

De Kogel, W. J., Van der Hoek, M., and Mollema, C. (1997b) Oviposition preference of western flower thrips for cucumber leaves from different positions along the plant stem. Entomol. Exp. Appl. 82:283-288.

De Jager, C. M., Butot, R. P. T., De Jong, T. J., Klinkhamer, P. G. L., and Van der Meijden, E. (1993) Population growth and survival of western flower thrips *Frankliniella occidentalis* Pergande (Thysanoptera: Thripidae) on different chrysanthemum cultivars: two methods for measuring resistance. J. Appl. Entomol. 115:519-525.

De Jager, C. M., Butot, R. P. T., Klinkhamer, P. G. L., De Jong, T. J., Wolff, K., and Van der Meijden, E. (1995) Genetic variation in chrysanthemum for resistance to *Frankliniella occidentalis* (Thysanoptera: Thripidae). Entomol. Exper. Appl. 77:277-287.

De Vries, E. J., Jacobs, G., and Breeuwer, J. A. J. (1995) Symbiotic bacteria of the western flower thrips, *Frankliniella occidentalis*. Proc. Exper. & Appl. Entomol. NEV 6:87-92.

De Vries, E. J., Breeuwer, J. A. J., Jacobs, G., and Mollema, C. (2001a) The association of western flower thrips, *Frankliniella occidentalis*, with a near *Erwinia* species gut bacteria: transient or permanent? J. Invertebr. Pathol. 77:120-128. .

De Vries, E. J., Jacobs, G., and Breeuwer, J. A. J. (2001b) Growth and transmission of gut bacteria in the Western Flower Thrips, *Frankliniella occidentalis*. J. Invertebr. Pathol. 77:129-137.

De Vries, E. J., Jacobs, G., Breeuwer, J. A. J., Sabelis, M. W., and Menken, S. B. J. (2004) Diet dependent fitness effects of gut bacteria on their insect host: the symbiosis of *Erwinia* sp. and western flower thrips. Proc. Royal Soc. Lond. B. 271:2171-2178.

De Vries, E. J., Vos, R. A., Jacobs, G., and Breeuwer, J. A. J. (2006) Western flower thrips (Thysanoptera: Thripidae) preference for thrips-damaged leaves over fresh leaves enable uptake of symbiotic gut bacteria. Eur. J. Entomol. 103:779-786.

De Vries, E. J., Van der Wurff, A., Jacobs, G., and Breeuwer, J. A. J. (2008) Onion thrips (*Thrips tabaci*) have gut bacteria that are closely related to the symbionts of western flower thrips (*Frankliniella occidentalis)*. J. Insect Science 23: 1-11.

Degnan, P. H., Lazarus, A. B., Brock, C. D., and Wernegreen, J. J. (2004) Host-symbiont stability and fast evolutionary rates in an ant-bacterium association: cospeciation of *Camponotus* species and their endosymbionts, *Candidatus* Blochmannia. Syst. Biol. 53:95-110.

Demaio, J., Pumpuni, C. B., Kent, M., and Beier, J. C. (1996) The midgut bacterial flora of wild *Aedes triseriatus*, *Culex pipiens*, and *Psorophora columbiae* mosquitoes. Am. J. Trop. Med. Hyg. 54:219-223.

Delalibera jr., I., Handelsman, J., and Raffa, K. F. (2005) Contrasts in cellulolytic activites of the gut microorganisms between the wood borer, *Saperda vestita* (Coleoptera: Cerambycidae), and the

bark beetles, *Ips pini* and *Dendroctonus frontalis* (Coleoptera: Curculionidae). Environ. Entomol. 34:541-547. .

Diehl, S. R., and Bush, G. L. (1984) An evolutionary and applied perspective of insect biotypes. Annu. Rev. Entomol. 29:471-504.

Dillon, R. J., and Charnley, A. K. (1995) Chemical barriers to gut infection in the desert locust: in vivo production of antimicrobial phenols associated with the bacteria *Pantoea agglomerans*. J. Invertebr. Pathol. 66:72-75.

Dillon, R. J., and Charnley, K. (1996) Colonization of the guts of germ-free desert locusts, *Schristocerca gregaria*, by the bacterium *Pantoea agglomerans*. J. Invertebr. Pathol. 67:11-14.

Dillon, R. J., Vennard, C. T., and Charnley, A. K. (2000) Exploitation of gut bacteria in the locust. Nature 403:851.

Dillon R. J., Vennard C. T., and Charnley A. K. (2002) A note: gut bacteria produce components of locust cohesion pheromone. J. Appl. Microbiol. 92:759-763.

Dillon, R. J., and Dillon, V. M. (2004) The gut bacteria of insects: nonpathogenic interactions. Annu. Rev. Entomol. 49:71-92.

Doebeli, M., and Knowlton, N. (1998) The evolution of interspecific mutualisms. Proc. Natl. Acad. Sci. USA 95:8676-8680.

Domingo, J. W. S., Kaufman, M. G., Klug, M. J., and Tiedje, J. M. (1998) Characterization of the cricket hindgut microbiota with fluorescent labeled rRNA-targeted oligonucleotide probes. Appl. Environ. Microbiol. 64:752-755.

Douglas, A. E. (1988) Experimental studies on the mycetome symbiosis in the leafhopper *Euscelis incisus*. J. Insect Physiol. 34:1043-1053.

Douglas, A. E. (1989) Mycetocyte symbiosis in insects. Biol. Rev. 64:409-434.

Douglas, A. E. (1992) Requirements of pea aphids (*Acyrtosiphon pisum*) for their symbiotic bacteria. Entomol. Exp. Appl. 65:195-198.

Douglas, A. E. (1994) Symbiotic Interactions. Oxford University Press, Oxford, UK. 148 pp.

Douglas, A. E. (1995) The ecology of symbiotic micro-organisms. Adv. Ecol. Res. 26:69-103.

Douglas, A. E. (1997) Provenance, experience and plant utilisation by the polyphagous aphid, *Aphis fabae*. Entomol. Exp. Appl. 83:161-170.

Douglas, A. E., and Smith, D. C. (1989) Are endosymbioses mutualistic? Trends Ecol. Evol. 4:350-352.

Douglas, A. E., and Beard, C. B. (1996) Microbial symbiosis in the midgut of insects. *In:* Biology of the Insect Midgut (M. J. Lehane and P. F. Billingsley, eds.), p. 419-431. Chapman and Hall, London, UK.

Drew, R. A. I., Courtice, A. C., and Teakle, D. S. (1983) Bacteria as a natural source of food for adult fruit flies (Diptera: Tephritidae). Oecologia 60:279-284.

Drew, R. A. I., and Lloyd, A. C. (1987) Relationship of fruit flies (Diptera: Tephritidae) and their bacteria to host plants. Ann. Entomol. Soc. Am. 80:629-636.

Dreyer, D. L., and Campbell, B. C. (1987) Chemical basis of host-plant resistance to aphids. Plant, Cell and Environment 10:353-361.

Dugas, J. E., Zurek, L., Paster, B. J., Keddie, B. A., and Leadbetter, E. R. (2001) Isolation and characterization of a *Chryseobacterium* strain from the gut of the American cockroach, *Periplaneta americana*. Arch. Microbiol. 175:259-262. .

Düggeli, M. (1904) Die Bakterienflora gesunder Samen und daraus gezogener Keimpflanzen. Zentralbl. Bakteriol. Parasitenk. Infektionskr. Abt II 12:602–614, and 13:56-63, and 13:198-207.

Dunn, A. K., and Stabb, E. V. (2005) Culture-independent characterization of the microbiota of the ant lion Myrmeleon mobilis (Neuroptera: Myrmeleontidae). Appl. Environ. Microbiol. 71:8784-8794.

Dye, D. W. (1964) The taxonomic position of *Xanthomonas trifolii* (Huss, 1907) James, 1955. New Zealand J. Sci. 7:261-269.

Dykes, G. A. (1995) Bacteriocins: ecological and evolutionary significance. Trends Ecol. Evol. 10:186-189.

Egert, M., Wagner, B., Lemke, Th., Brune, A., and Friedrich, M. W. (2003) Microbial community structure in midgut and hindgut of the humus-feediong larva of *Pachnoda ephippiata* (Coleoptera: Scarabeidae). Appl. Environ. Microbiol. 69:6659-6668.

Eigenbrode, S. D., Ding, H. J., Shiel, P., and Berger, P. H. (2002) Volatiles from potato plants infected with potato leafroll virus attract and arrest the virus vector, *Myzus persicae* (Homoptera: Aphididae). Proc. Royal Soc. Lond. B. 269:455-460.

Eisenbach, J., and Mittler, T. E. (1987) Extra-nuclear inheritance in sexually produced aphid: the ability to overcome host plant resistance by biotype hybrids of the greenbug, *Schizaphis graminum.* Experientia 43:332-334.

Elliot, C., and Poos, T. W. (1940) Seasonal development of insect vectors and host range of bacterial wilt on sweet corn. J. Agricult. Res. 60:645-686.

Endress, P. K. (1994) Diversity and Evolutionary Biology of Tropical Flowers. Cambridge Univ. press, Cambridge, UK. 354 pp.

Espinoza-Fuentes, F. P., and Terra, A. R. (1987) Physiological adaptations for digestion bacteria. Water fluxes and distribution of digestive enzymes in *Musca domestica* larval midgut. Insect Biochem. 17:809-817.

Eutick, M. L., Veivers, P., O'Brien, R. W., and Slaytor, M. (1987) Dependence of the higher termite, *Nasutitermes exitiosus* and the lower termite, *Coptotermes lacteus* on their gut flora. J. Insect Physiol. 24:383-368.

Ewald, P. (1994) Evolution of infectious disease. Oxford University Press, Oxford, UK.

Ewing, W. H., and Fife, M. A. (1972) *Enterobacter agglomerans* (Beijerinck) comb. nov. (the Herbicola-Lathyri Bacteria). Int. J. Syst. Bacteriol. 22:4-11.

Farrar, J. J., and Davis, R. M. (1991) Relationships among ear morphology, western flower thrips and *Fusarium* ear rot of corn. Phytopathology 81:661-666.

Fitt, G. P., and O'Brien, R. W. (1995) Bacteria associated with four species of *Dacus* (Diptera: Tephritidae) and their role in the nutrition of larvae. Oecologia 67:447-454.

Fery, R. L., and Schalke, J. M. (1991) Resistance in pepper (*Capsicum annuum* L.) to western flower thrips (*Frankliniella occidentalis* [Pergande]). Hort. Science 26:1073-1074.

Fox, L. R., and Morrow, P. A. (1981) Specialization: species property or local phenomenon? Science 211:887-893.

Franssen, C. J. H., and Mantel, W.P. (1962) Lijst van in Nederland aangetroffen Thysanoptera met beknopte aantekeningen over hun levenswijze en hun betekenis voor onze cultuurgewassen. *Tijdschr. Entomol.*, 105:97-133.

Frey, J. E. (1993) The analysis of arthropod pest movement through trade in ornamental plants. BCPC Monograph 54:157-165.

Frey, J. E., and Frey, B. (2004) Origin of intra-individual variation in PCR-amplified mitochondrial cytochrome oxidase I of *Thrips tabaci* (Thysanoptera: Thripidae): mitochondrial heteroplasmy or nuclear integration? Hereditas 140:92-98. .

Fukatsu, T., and Nikoh, N. (1998) Two intracellular symbiotic bacteria from the mulberry psyllid *Anomoneura mori* (Insecta, Homoptera). Appl. Environ. Microbiol. 64:3599-3606.

Fukatsu, T., and Nikoh, N. (2000a) Endosymbiotic microbiota of the bamboo pseudococcid *Antonina crawii* (Insecta, Homoptera). Appl. Environ. Microbiol. 66:643-650. .

Fukatsu, T., Nikoh, N., Kawai, R., and Koga, R. (2000b) The secondary endosymbiotic bacterium of the pea aphid *Acyrtosiphon pisum* (Insecta: Homoptera). Appl. Environ. Microbiol. 66:2748-2758.

Gaston, K. J., and Mound, L. A. (1983) Taxonomy, hypothesis testing, and the biodiversity crisis. Proc. Roy. Soc. London (B) 251:139-142.

Gaum, W. G., Giliomee, J. H., and Pringle, K. L. (1994) Life history and life tables of western flower thrips, *Frankliniella occidentalis* (Thysanoptera: Thripidae), on English cucumbers. Bull. Entomol. Res. 84:219-224.

Gavini, F., Mergaert, J., Beji, A., Mielcarek, C., Izard, D., Kersters, K., and De Ley, J. (1989) Transfer of *Enterobacter agglomerans* (Beijerinck, 1888) Ewing and Fife 1972 to *Pantoea* gen. nov. as *Pantoea agglomerans* comb. nov. and description of *Pantoea dispersa* sp. nov. Int. J. Syst. Bacteriol. 39:337-345.

Genkai-Kato, M., and Yamamura, N. (1999) Evolution of symbiosis without vertical transmission. Theor. Pop. Biol. 55:309-323.

Genta, F. A., Dillon, R. J., Terra, W. R., and Ferreira, C. (2006) Potential role for gut microbiota in cell wall digestion and glucoside detoxification in *Tenebrio molitor* larvae. J. Insect Physiol. 52:593-601.

Giliomee, J. H. (1989) First record of western flower thrips, *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) from South Africa. J. Entomol. Soc. Sth. Afr. 52:179-180.

Gilliam, M. (1997) Identification and roles of non-pathogenic microflora associated with honey bees. FEMS Microbiol Ecol. 155:1-10.

Gilliam, M., and Valentine, D. K. (1974) Enterobacteriaceae isolated from foraging worker honey bees, *Apis mellifera.* J. Invertebr. Pathol. 23:38-41.

Gilliam, M., and Lohrenz, B. J. (1983) Gram-positive cocci from apiarian sources. J. Invertebr. Pathol. 42:187-195.

Gillings, M. R., Rae, D., Heron, E. A., and Beattie, G. A. C. (1995) Tracking thrips populations with DNA-based methods. Proc. 1995 Austr. New-Zl. Thrips workshop. NSW Agricult. Gosford, p. 47-103.

Gitaitis, R. D., Walcott, R. R., Wells, M. L., Perez, J. C. D., and Sanders, F. H. (2003) Transmission of *Pantoea ananatis*, causal agent of center rot of onion, by tobacco thrips, *Frankliniella fusca*. Plant Disease 87:675-678.

Goldbach, R. W, and Peters, D, (1994) Possible causes of the emergence of tospovirus diseases. Seminars of Virology 5:113-120.

Goodchild, A. J. P. (1955) The bacteria associated with *Triatoma infestans* and some other species of Reduviidae. Parasitology 54:441-448.

Gould, F. (1979) Rapid host range evolution in a population of the phytophagous mite *Tetranychus urticae* Koch. Evolution 33:791-802.

Grassé, P.-P., and Noirot, Ch. (1959) L'évolution de la symbiose chez les Isoptères. Experientia 15:365-372.

Greenberg, B. (1968) Model for the deconstruction of bacteria in the midgut of blow-fly maggots. J. Med. Entomol 5:31-38. .

Grisham, M. P., Sterling, W. L., Powell, R. D., and Morgan, P. W. (1987) Characterization of the induction of stress ethylene synthesis in cotton caused by the cotton fleahopper (Hemiptera: Miridae) and its microorganisms. Ann. Entomol. Soc. Am. 80:411-416.

Guldemond, J. A., Tigges, W. T., and De Vrijer, P. W. F. (1994) Host races of *Aphis gossypii* (Homoptera: Aphididae) on cucumber and chrysanthemum. Environ. Entomol. 23:1235-1240.

Gijzen, H. J., Van der Drift, Chr., Barugahare, M., and Op den Camp, H. J. M. (1994) Effect of host diet and hindgut microbial composition on cellulolytic activity in the hindgut of the American cockroach, *Periplaneta americana*. Appl. Environ. Microbiol. 60:1822-1826.

Haas, F., and König, H. (1987) Characterisation of an anaerobic symbiont and the associated aerobic bacterial flora of *Pyrrhocorus apterus* (Heteroptera: Pyrrhocoridae). FEMS Microbiol. Ecol. 45:99-106.

Hadrys, H., Balick, M., and Schierwater, B. (1992) Applications of random amplified polymorphic DNA (RAPD) in molecular ecology. Mol. Ecol. 1:55-63.

Hagen, K. S. (1966) Dependence of the olive fly, *Dacus oleae*, larvae on symbiosis with *Pseudomonas savastanoi* for the utilization of olive. Nature 209:423-424.

Harada, H., Oyaizu, H., and Ishikawa, H. (1996) A consideration about the origin of aphid intracellular symbiont in connection with gut bacterial flora. J. Gen. Appl. Microbiol. 42:17-26.

Harley, J. E., and Smith, S. E. (1983) Mycchorizal Symbiosis. Academic Press, London, UK. 530 pp.

Harrewijn, P., Tjallingii, F., and Mollema, C. (1996). Electrical recording of plant penetration by western flower thrips. Entomol. Exp. Appl. 79:345-353.

Hauben, L., Moore, E. R. B., Vauterin, L., Steenackers, M., Mergaert, J., Verdonck, L., and Swings, J. (1998) Phylogenetic position of phytopathogens within the Enterobacteriaceae. System. Appl. Microbiol. 21:384-397.

Haynes, S., Darby, A. C., Daniell, T. J., Webster ,G. Van Veen, F. J. F., Godfray, H. C. J., Prosser, J. I., and Douglas, A. E. (2003) Diversity of bacteria associated with natural aphid populations. Appl. Environ. Microbiol. 69:7216-7223.

Heddi, A., Grenier, A.-M., Khatchadourian, C., Charles, H., and Nardon, P. (1999) Four intracellular genomes direct Weevil biology: nuclear, mitochondrial, principal endosybiont, and *Wolbachia*. Proc. Natl. Acad. Sci. USA 96:6814-6819.

Helyer, N. L., and Brobyn, P. J. (1992) Chemical control of western flower thrips (*Frankliniella occidentalis* Pergande). Ann. Appl. Biol. 121:219-231.

Herre, E. A., Knowlton, N., Mueller, U. G., and Rehner, S. A. (1999) The evolution of mutualisms: exploring the paths between conflict and cooperation. Trends Evol. Evol. 14:49-53.

Hill, P., Campbell, J. A., and Petrie, I. A. (1976). *Rhodnius prolixus* and its symbiotic actinomycete: a microbiological, physiological and behavioural study. Proc. R. Soc. Lond. B. 194:501-524.

Hirose, E., Panizzi, A. R., De Souza, J. T., Cattelan, A. J., and Aldrich, J. R. (2006) Bacteria in the gut of souther green stink bug (Heteroptera: Pentatomidae). Ann. Entomol. Soc. Am. 99:91-95.

Hochberg, M. E., Gomulkiewicz, R., Holt, R. D., and Thompson, J. N. (2000) Weak sinks could cradle mutualistic symbiosis – strong sources should harbour parasitic symbiosis. J. Evol. Biol. 13:213-222.

Hochberg, M. E., and Holt, R. J. (2002) *In:* The Adaptive Dynamics of Infectious Diseases: In Pursuit of Virulence Management (U. Dieckmann, J. A .J. Metz, M. W. Sabelis and K. Sigmund eds.) Cambridge University Press, Cambridge, UK, .

Holden, P. R., Brookfield, J. F. Y., and Jones, P. (1993) Cloning and characterization of an *ftsZ* homologue from a bacterial symbiont of *Drosophila melanogaster*. Mol. Gen. Genet. 240:213-230.

Hormaeche, E., and Edwards, P. R. (1960) A proposed genus *Enterobacter*. International Bulletin of Bacteriological Nomenclature and Taxonomy. 10:71-74.

Hough, J. A., Harman, G. E., and Eckenrode, C. J. (1981) Microbial stimulation of onion maggot oviposition. Environ. Entomol. 10:206-211.

Houk, E. J., and Griffiths, G. W. (1980) Intracellular symbiotes of the Homoptera. Annu. Rev. Entomol. 25:161-187.

Howard, D. J., Bush, G. L., and Breznak, J. A. (1985) The evolutionary significance of bacteria associated with *Rhagoletis*. Evolution 39:405-417.

Hoyt, C. P., and Osborne, G. O. (1971) Production of an insect sex attractant by symbiotic bacteria. Nature 230:472-473.

Humphreys, N. J., and Douglas, A. E. (1997) Partitioning of symbiotic bacteria between generations of an insect: a quantitative study of a *Buchnera* sp. in the Pea Aphid (*Acyrtosiphon pisum*) reared at different temperatures. Appl. Environ. Microb. 63:3294-3296.

Hunt, J., and Charnley, A. K. (1981) Abundance and distribution of the gut flora of the desert locust *Schristocerca gregaria*. J. Invertebr. Pathol. 38:378-385.

Hunter, W. B., and Ullman, D. E. (1992) Anatomy and ultrastructure of the piercing sucking mouthparts and paraglossal sensilla of *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae). Int. J. Insect Morphol. Embryol. 18:161-171.

Hurst, L. D. (1993) The incidence, mechanisms and evolution of cytoplasmic sex ratio distorters in animals. Biol. Rev. 68:121-193.

Husseneder, C., and Grace, J. K. (2005) Genetically engineered termite gut bacteria *(Entrobacter cloacae*) deliver and spread foreign genes in termite colonies. Appl. Microbiol. .

Hypsa, V. (1993) Endocytobionts of *Triatoma infestans*: distribution and transmission. J. Invertebr. Pathol. 61:32-38.

Hypsa, V., and Dale, C. (1997) In vitro culture and phylogenetic analysis of "*Candidatus* Arsenophonus triatominarum", an intracellular bacterium from the Triatomine bug, *Triatoma infestans*. Int. J. Syst. Bacteriol. 47:1140-1144.

Immaraju, J. A., Paine, T. D., Bethke, J. A., Robb, K. L., and Newman, J. P. (1992) Western flower thrips (Thysanoptera: Thripidae) resistance to insecticides in coastal California greenhouses. J. Econ. Entomol. 85:9-14.

Ishikawa, H. (1981) Biochemical and molecular aspects of endosymbiosis in insects. Int. Rev. Cytol. 116:1-43.

Iverson, K. L., Bromel, M. C., Anderson, A. W., and Freeman, T. P. (1984) Bacterial symbionts in the sugar beet maggot, *Tetanops myopaeformis* (Von Röder). Appl. Environ. Microbiol. 47:22-33.

Jacobsen, R. (1997) Integrated pest management (IPM) in glasshouses. *In:* Thrips as Crop Pest (T. Lewis, ed.), p. 639-666. CAB Int., New York, NY, USA.

Jaenike, J. (1990) Host specialisation in phytophagous insects. Annu. Rev. Ecol. Syst. 21:243-273.

Janssen, A., Pallini, A., Venzon, M., and Sabelis, M. W. (1998) Behaviour and indirect interactions in food webs of plant-inhabiting arthropods. Exper. Appl. Acarol. 22:497-521.

Jarosz, J. (1979) Gut flora of *Galleria melonella* suppresing ingested bacteria. J. Invertebr. Pathol. 34:192-198.

Jones, C.G. (1984) Micro-organisms as mediators of plant resource exploitation by insect herbivores. *In:* A New Ecology, Novel Approaches to Interactive Systems (E. A. Bernays, ed.), pp. 53-98. John Wiley and sons, New York (N.Y.), USA.

Jongsma, M. A. (1995) The resistance of insects to plant proteinase inhibitors. Ph. D. thesis, Landbouwuniversiteit Wageningen, 93 pp.

Kadavy, D. R., Plantz, B., Shaw, Chr. A., Myatt, J., Kokjohn T. A., and Nickerson, K. W. (1999) Microbiology of the oil fly, *Helaeomyia petrolei*. Appl. Environ. Microbiol. 65:1477-1482.

Kaltenpoth, M., Göttler, W., Herzner, G., and Strohm, E. (2005) Symbiotic bacteria protect wasp larvae from fungal infestation. Curr. Biol. 15:475-479.

Kaneko, J., Yoshida, T., Owada, T., Kita, K., and Tanno, K. (1995) *Erwinia herbicola:* Ice-nucleation active bacteria isolated from the diamondback moth, *Plutella xylostella* L. pupae (in Japanese). Jpn. J. Appl. Entomol. Zool. 35:247-251.

Karban, R. (1989) Fine-scale adaptation of herbivorous thrips to individual host plants. Nature 340:60-61.

Kaufman, M. G., Klug, M. J., and Merrit, R. W. (1989) Growth and food utilization parameters of germ-free house crickets, *Acheta domesticus*. J. Insect Physiol. 35:957-967.

Kaufman, M. G., and Klug, M. J. (1991) The contribution of hindgut bacteria to dietary carbohydrate utilization by crickets (Orthoptera: Gryllidae). Comp. Biochem. Physiol. 98A:117-123.

Keddy, P. (1990) Is mutualism really irrelevant to ecology? Biol. Ecol. Soc. Amer. 71:101-102.

Kikkert, M., Meurs, C., Van de Wetering, F., Dorfmüller, S., Peters, D., Kormelink, R., and Goldbach, R.W., (1998) Binding of Tomato Spotted Wilt Virus to a 94-kDa thrips protein. Phytopathology 88:63-69.

Kirk, W. D. J. (1984) Pollen-feeding in thrips (Insecta: Thysanoptera). J. Zool. 204:107-117.

Kirk, W. D. J. (1985a) Pollen-feeding and the host specificity and fecundity of flower thrips (Thysanoptera). Ecol. Entomol. 10:281-289.

Kirk, W. D. J. (1985b) Aggregation and mating of thrips in flowers of *Calystega sepium*. Ecol. Entomol. 10:433-440.

Kirk, W. D. J. (1988) Thrips and pollination biology. *In:* Dynamics of Insect-Plant Interaction Recent Advances and Future Trends (T. N. Ananthakrishan, and A. Raman, eds.), p. 124-135. Oxford and IBH, New Delhi, India. .

Kirk, W. D. J. (1997a) Feeding. *In:* Thrips as Crop Pest (T. Lewis, ed.), CAB Int., New York, NY, USA.

Kirk, W. D. J. (1997b) Distribution, abundance and population dynamics. *In:* Thrips as Crop Pest (T. Lewis, ed.), p. 217-258. CAB Int., New York, NY, USA.

Klug, M. J., and Kotarski, S. (1980) Bacteria associated with the gut tract of larval stages of the aquatic cranefly *Tipula abdominalis* (Diptera: Tipulidae). Appl. Environ. Microbiol. 40:408-416.

Koch, A. (1967) Insects and their endosymbionts. *In:* Symbiosis. Volume II; Associations of Invertebrates, Birds, Ruminants and Other Biota (S. Mark Henry, ed.), p. 1-106. Acad. Press, New York, NY, USA.

Kumar, N. K. K., Ullman, D. E., and Cho, J. J. (1995) Resistance among *Lycopersicon* species to *Frankliniella occidentalis* (Thysanoptera: Thripidae). J. Econ. Entomol 88:1057:1065.

Kuzina, L. V., Miller, E. D., Ge, B., and Miller, T. A. (2002) Transformation of *Erwinia gergoviae* isolated from the pink bollworm (Lepidoptera: Gelechiidae) gut with *Bacillus thuringiensis* toxin. Curr. Microbiol. 44:1-4.

Lane, D. J. (1991) 16S/23S rDNA sequencing. *In:* Sequencing and Hybridization Techniques in Bacterial Systematics (E. Stackebrandt and M. Goodfellow, eds.), pp. 230-270. J. Wiley and sons, Chichester, UK. .

Lauzon, C. R., Sjogren, R. E., and Prokopy, R. J. (2000) Enzymatic capabilities of bacteria associated with apple maggot flies: a postulated role in attration. J. Chem. Ecol. 26:953-967.

Law. R., and Diekmann, U. (1998) Symbiosis through exploitation and the merger of lineages in evolution. Proc. R. Soc. Lond. B. 265:1245-1253.

Leigh, T. F. (1995) Bionomics of cotton thrips. *In:* Thrips Biology and Pest Management (B. L. Parker, ed.), p. 61-70. Plenum Press, New York, NY, USA.

Lefevre, C., Charles, H., Vallier, A., Delobel, B., Farrell, B., and Heddi, A. (2004) Endosymbiont phylogenesis in the Dryophthoridae Weevils: evidence for bacterial replacement. Mol. Biol. Evol. 21:965973.

Lewis, T. (1997) Thrips as Crop Pest .CAB Int., New York, NY, USA. 594 pp.

Lind, E., and Uring, J. (1986) Clinical strains of *Enterobacter agglomerans* (synonyms: *Erwinia herbicola*, *Erwinia milletiae*) identified by DNA-DNA hybridization. Acta Path. Immunol. Scand. Sect. B 94:205-213.

Lilley, A. K., Hails, R. S., Cory, J. S., and Bailey, M. J. (1997) The dispersal and establishment of pseudomonad populations in the phyllosphere of sugar beet by phytophagous caterpillars. FEMS Microbiol. Ecol. 24:151-157.

Liu, W., Zhao, G., and Knowles, C. O. (1994) Substrate specificity and inhibitor sensitivity of cholinesterases in homogenates of western flower thrips. Pesticid. Biochem. Physiol. 49:121-131.

Lloyd, A. C., Drew, R. A. L., Teakle, D. S., and Hayward, A. C. (1986) Bacteria associated with some *Dacus* species (Diptera: Tephritidae) and their host fruit in Queensland. Aust. J. Biol. Sci. 39:361-368.

Lublinkhof, J., and Foster, D. E. (1977) Development and reproductive capacity of *Frankliniella occidentalis* (Thysanoptera: Thripidae) reared at three temperatures. J. Kansas Entomol. Soc. 50:313-316.

Lukwinski, A. T., Hill, J. E., Khatchatourians, G. G., Hemmingsen, S. M., and Hegedus, D. D. (2006) Biochemical and taxonomic characterisation of bacteria associated with the crucifer root maggot (*Delia radicum*). Can. J. Microbiol. 52:197-208.

Maccollom, G. B., Lauzon, C. R., Weires, R. W., and Rutkowski, A. A. (1992) Attraction of adult apple maggot (Diptera, Tephritidae) to microbial isolates. J. Econ. Entomol. 85:83-87.

MacCollom, G. B., Lauzon, C. R., Payne, E. B., and Currier, W. W. (1994) Apple maggot (Diptera: Tephritidae) trap enhancement with washed bacterial cells. Environ. Entomol. 23:354-359.

Mantel, W. P., and Van de Vrie, M. (1988) De Californische trips, *Frankliniella occidentalis*, een nieuwe schadelijke tripssoort in de tuinbouw onder glas in Nederland. Entomol. Ber. 48:140-144.

Margulis, L. (1993) Symbiosis in Cell Evolution: Microbial Communities in the Archaen and Protozoic Eons (2nd ed.). W.H. Freeman and co., New York, NY, USA. 452 pp.

Maris, P. C., Joosten, N. N., Goldbach, R. W., and Peters, D. (2004) Tomato spotted wilt virus infection improves host suitability for its vector *Frankliniella occidentalis*. Phytopathol. 94:706-711.

Marshall, S. A., and Eymann, M. (1981) Micro-organisms as food for the onion maggot, *Delia antiqua* (Diptera: Anthomyiidae). Proc. Entomol. Soc. Ontario 112:1-5.

Martin, N. A., and Workman, P. J. (1994) Confirmation of a pesticide-resistant strain of western flower thrips in New Zealand. Proc. 47th New Zealand Plant Prot. Conf. p. 144-147.

Martinez, A. J., Robacker, D. C., Garcia, J. A., and Esau, K. L. (1994) Laboratory and field olfactory attraction of the Mexican fruit fly (Diptera: Tephritidae) to metabolites of bacterial species. Florida Entomologist 77:117-126.

Matteson, N., and Terry, L. I. (1992) Response to color of male and female *Frankliniella occidentalis* during swarming and non-swarming behavior. Entomol. Exp. Appl. 63:187-201.

Maynard-Smith, J., and Szathmáry, E. (1995) The Major Transitions in Evolution. Freeman and co., Oxford, UK. 346 pp.

McDonald, O. (1995) The response of western flower thrips to dichlorvos and malathion in the United Kingdom. *In:* Thrips Biology and Pest Management (B. L. Parker, M. Skinner, and T. Lewis, eds.), p. 347-350. Plenum Press, New York, NY, USA.

McKenzie, A. (1996) A trade-off for host plant utilization in the black bean aphid, *Aphis fabae*. Evolution 50:155-162.

Mead, L. J., Khachatourians, G. G., and Jones, G. A. (1988) Microbial ecology of the gut in laboratory stocks of the migratory grasshopper, *Melanoplus sanguinipes* (Fab.) (Orthoptera: Acrididae). Appl. Environ. Microbiol. 54:1174-1181.

Menken, S. B. J. (1996) Pattern and process in the evolution of insect-plant associations: *Yponomeuta* as an example. Entomol. Exp. Appl. 80:297-305.

Mergaert, J., Verdonck, L., Kersters, K., Swings, J., Boeufgras, J.-M., and De Ley, J. (1984) Numerical taxonomy of *Erwinia* species using API systems. J. Gen. Microbiol. 130:1893-1910.

Mergaert, J., Verdonck, L. and Kesters., K. (1993) Transfer of *Erwinia ananas* (synonym, *Erwinia uredovera* ) and *Erwinia stewartii* to the genus *Pantoea* emend. as *Pantoea ananas* (Serrano 1928) comb. nov. and *Pantoea stewartii* (Smith, 1898) comb.nov., respectively, and description of *Pantoea stewartii* subsp. *indologenes* subsp. nov. Int. J. Syst. Bacteriol. 43:162-173.

Mollema, C., Steenhuis, G., and Inggamer, H. (1995). Genotypic effects of cucumber responses to infestation of western flower thrips. *In:* Thrips Biology and Pest Management (B. L. Parker, M. Skinner, and T. Lewis, eds.), p. 397-402. Plenum Press, New York, NY, USA. .

Moore, G. E. (1972) Microflora from the alimentary tract of healthy southern pine beetles, *Dendroctonus frontalis* (Scolytidae), and their possible relationship to pathogenicity. J. Invertebr. Pathol. 19:72-75.

Moran, N., and Baumann, P. (1994) Phylogenetics of cytoplasmically inherited microorganisms of arthropods. Trends Ecol. Evol. 9:15-20.

Moran, N. A., Russell, J. A., Koga, R., and Fukatsu, T. (2005) Evolutionary relationshps of three new species of *Enterobacteriaceae* living as symbionts of aphids and other insects. Appl. Environ. Microbiol. 71:3302-3310.

Moritz, G. (1996) Ontogenic development of the digestive systems of thrips. *In:* Proc. of the XX International Congres of Entomology, p. 458. Firenze, Italia.

Moritz, G. (1997) Structure, growth and development. *In:* Thrips as Crop Pest (T. Lewis, ed.). CAB Int., New York, NY, USA. .

Morse, J. G., and Hoddle, M. S. (2006) Invasion biology of thrips. Annu. Rev. Entomol. 51:67-89.

Mound, L. A. (1997) Biological diversity. *In:* Thrips as Crop Pest (T. Lewis, ed.). CAB Int., New York, NY, USA.

Mound, L. A., Heming, B. S., and Palmer, J. M. (1980) Phylogenetic relationships between the families of recent Thysanoptera. Zool. J. Linnean Soc. London 69:111-141.

Mound, L. A., and Teulon, D. A. J. (1995) Thysanoptera as phytophagous opportunists. *In:* Thrips Biology and Pest Management (B. L. Parker, M. Skinner, and T. Lewis (eds.), p. 3-19. Proc. of the 1993 International Conference on Thysanoptera, 28-30 Sept. 1993. Plenum press, New York, NY, USA.

Mound, L., Marullo, R., Goldarenza Lafuente, A., Loomans, A. J. M., Moritz, G., Ready. P. D., and Testa, J. M. (1998) Pest Thysanoptera of mediterranean fruit flies, final report. 12 pp.

Munson, M. A., Baumann, P., and Kinsey, M. G. (1991) *Buchnera* gen. nov. and *Buchnera aphidicola*, sp. nov., a taxon consisting of the mycetocyte-associated, primary endosymbionts of aphids. Int. J. System. Bacteriol. 41:566-568.

Munson, M. A., Baumann, P., and Moran, N. A. (1992) Phylogenetic relationships of the endosymbiosis of mealybugs (Homoptera: Pseudococcidae) based on 16S rDNA sequences. Molec. Phyolgen. Evol. 1:26-30.

Murai, T., and Ishii, T. (1982) Simple rearing methods for flower thrips (Thysanoptera: Thripidae) on pollens (in Japanese). Jpn. J. Appl. Entomol. Zool. 26:149-154.

Murai, T. (1990) Parthenogenic reproduction in *Thrips tabaci* and *Frankliniella intonsa* (Insecta: Thysanoptera). Adv. Invertebr. Repr. 5:357-362.

Murphy, K. M., Teakle, D. S., and McRae, I. C. (1994) Kinetics of colonization of adult Queensland Fruit flies (*Bactrocera tryoni*) by dinitrogen-fixing alimentary tract bacteria. Appl. Environ. Microbiol. 60:2508-2517.

Musgrave, A. J. (1964) Insect mycetomes. Can. J. Entomol. 1:377-389.

Nagata, T. (1999) Competence and Specificity of Thrips in the Transmission of Tomato Spotted Wilt Virus. Ph D thesis, Wageningen Agricultural University, 96 pp.

Nagata, T., Inoue-Nagata, A. K., Smid, H. M., Goldbach, R. W., and Peters, D., (1999) Tissue tropism related to vector competence of *Frankliniella occidentalis* for tomato spotted wilt tospovirus. J. Gen. Virol. 80:507-515.

Nagelkerke, C. (1996) Evolution of Sex-allocation Strategies in Predatory Mites, Ph D thesis Univ. of Amsterdam. 227 pp. .

Nardi, J. B., Bee, Ch. M., Miller, L. A., Nguyen, N. H., Suh, S.-O., and Blackwell, M. (2006) Communities of microbes that inhabit the changing hindgut landscape of a subsocial beetle. Arthr. Struct. Develop. 35:57-68.

Nardon, P. (1973) Obtention d'une souche asymbiotique chez le charancon *Sitophilus sasakii* Tak.: differentes méthodes et comparaison avec la souche asymbiotique d'origine. CR Acad. Sci. 277:981-984.

Nardon, P., Wicker, C., Grenier, A.-M., et Laviolette, P. (1978) Contrôle de la prolifération des symbiotes par l'hôte: étude préliminaire de l'exo-b-N-acétylglucosaminidase chez le curculionide *Sitophilus oryzae* L. C. R. Acad. Sci. Paris 287:1157-1160.

Nardon, P., and Grenier, A. M. (1988) Genetical and biochemical interactions between the host and its endocytobionts in the weevils *Sitophilus* (Coleoptere, Curculionidae) and other related species. NATO ASI Series H 17:255-270.

Nault, B. A., Shelton, A. M., Gangloff-Kaufmann, J. L., Clark, M. E., Werren, J. H., Cabrera-La Rosa, J. C., and Kennedy, G.G. (2006) Reproductive modes in onion thrips (Thysanoptera: Thripidae) populations from New York onion fields. Environ. Entomol. 35:1663-1271.

Nei, M. (1987) Molecular Evolutionary Genetics. Columbia University press, New York, NY, USA. .

Nogge, G. (1978) Aposymbiotic tsetse flies, *Glossina morsitans morsitans*, obtained by feeding on rabbits immunized specifcally with symbionts. J. Insect Physiol. 24:290-304.

Nogge, G., and Gerresheim, A. (1982) Experiments on the elimination of symbionts from the tsetse fly, *Glossina morsitans morsitans* (Diptera: Glossinidae), by antibiotics and lysozyme. J. Invertebr. Pathol. 40:166-179.

O'Neill, S. L., Gooding, R. H., and Aksoy, S. (1993) Phylogentically distant symbiotic microorganisms reside in *Glossina* midgut and ovary tissues. Med. Veter. Entomol. 7:377-383.

Oakey, H. J., Ellis, J. T., and Gibson, L. F. (1996) Differentiation of *Aeromonas* genomospecies using random amplified polymorphic DNA polymerase chain reaction. J. Appl. Bact. 80:402-410.

Odinot, P. T., Meis, J. F. G. M., van den Hurk, P. J. J. C., Hoogkamp-Kastanje, J. A. A., and Welchers, W. J. G. (1995) PCR-based characterization of *Yersinia enterocolitica*: comparison with biotyping and serotyping. Epidemiol. Infect. 115:269-277.

Ohkuma, M., and Kudo, T. (1996) Phylogenetic diversity of the intestinal bacterial community in the termite *Reticulitermes speratus*. Appl. Environ. Microbiol. 62:461-468.

Olsen, G. J., and Woese, C. R. (1993) Ribosomal RNA: a key to phylogeny. FASEB J. 7:1113-1123.

Pallini Filho, A. (1999) Odour-mediated indirect interactions in an arthropod food web. Ph D-thesis, Univ. of Amsterdam. 91 pp. .

Pallini, A., Janssen, A., and Sabelis, M. W. (1997) Odour-mediated responses of phytophagous mites to conspecific and heterospecific competitors. Oecologia 110:179-185.

Parent, J.-G., Lacroix, M., Pagé, D., and L. Vézina. (1996) Identification of *Erwinia carotovera* from soft root diseased plants by Random Amplified Polymorphic DNA (RAPD) analysis. Plant Dis. 80:494-499.

Petri, L. (1904) Sopra la particolare localizzazione di una colonia batteria nel tubo diregente della larva della mosca olearia. Atti Reale Acad. Lincei [5] Classe Sci. Fis. Mat. e Nat. 13:506.

Peloquin, J. J., and Greenberg, L. (2003) Indentification of midgut bacteria from fourth instar red imported fire ant larvae, *Solenopsis invicta* Buren (Hymenoptera: Formicidae). J. Agricult. Urban Entomol. 20:157-164.

Pickett, C. H., Wilson, L. J., and Gonzales, D. (1988) Population dynamics and within-plant distribution of the western flower thrips [Thysanoptera: Thripidae] an early predator of spider mites infesting cotton. Environ. Entomol. 17:551-559.

Pierre, L. L. (1964) Uricase activity of isolated symbionts and the aposymbiotic fat body of a cockroach. Nature 201:54-55.

Potrikus, C. J., and Breznak, J. A. (1977) Nitrogen-fixing *Enterobacter agglomerans* isolated from guts of wood-eating termites. Appl. Environ. Microbiol. 33:392-399.

Ragsdale, D. W., Larson, A. D., and Newsom, L. D. (1979) Microorganisms associated with feeding and from various organs of *Nezara viridula*. J. Econ. Entomol. 72:725-727.

Resende, R. de O., De Avila, A. C., Goldbach, R. W., and Peters, D. (1991) Comparison of polyclonal antisera in the detection of tomato spotted wilt virus using double antibody sandwich and cocktail ELISA. J. Pathol. 132:46-56.

Robb, K. L. (1989) Analysis of *Frankliniella occidentalis* (Pergande) as a Pest of Floricultural Crops in Californian Greenhouses. Ph D thesis, Univ. of California, Riverside. 135 pp.

Roselló-Mora, R., and Amman, R. (2001) The species concept for prokaryotes. FEMS Microbiol. Rev. 25:39-67.

Rossiter, M. C. (1987) Genetic and phenotypic variation in diet breadth in a generalist herbivore. Evol. Ecol. 1:272-282.

Rubio, R. E. P., and McFadden, M. W. (1966) Isolation and identification of bacteria in the digestive tract of the Mexican fruit fly, *Anastrepha ludens* (Diptera: Tephritidae). Ann. Entomol. Soc. Am. 59:1015-1016.

Sabelis, M. W., and Van Rijn, P. C. J. (1997) Predation by insects and mites. *In:* Thrips as Crop Pest (T. Lewis, ed.), p. 259-354. CAB Int., New York, N.Y., USA.

Sasaki, T., Hayashi, H., and Ishikawa, H. (1991) Growth and reproduction of the symbiotic and aposymbiotic pea aphids, *Acyrtosiphon pisum*, maintained on artificial diets. J. Insect Physiol. 37:749-756.

Sauer, Chr., Stackebrandt, E., Gadau, J., Hölldobler, B., and Gross, R. (2000) Systematic relatonships and cospeciation of the bacterial endosymbionts and their Carpenter Ant host species: proposal of the new taxon *Candidatus* Blochmannia gen. nov. Int. J. Syst. Evol. Microbiol. 50:1877-1886. .

Saxena, R. C., and Barrion, A. A. (1987) Biotypes of insect pests of agricultural crops. Ins. Sci. Appl. 8:453-458.

Schmitt-Wagner, D., Friedrich, M. W., Wagner, B., and Brune, A. (2003) Phylogenetic diversity, abun-

dance, and axial distribution of bacteria in the intestinal tract of two soil-feeding termites (*Cubitermes* spp.). Appl. Environ. Microbiol. 69:6007-6017.

Schröder, D., Deppisch, H., Obermayer, M., Krohne, G., Stackebrandt, E., Hölldobler, B, Goebel, W., and Gross, R. (1996) Intracellular endosymbiotic bacteria of *Camponotus* species (carpenter ants): systematics, evolution and ultrastructure characterization. Molec. Microbiol. 21:479-489.

Schultz, J. E., and Breznak, J. A. (1978) Heterotrophic bacteria present in the hindguts of wood-eating termites [*Recticulitermes flavipes* (Kollar)]. Appl. Environ. Microbiol. 35:930-936.

Sherratt, T. N., and Roberts, G. (2002) The stability of co-operation involving variable investment. J. Theor. Biol. 215 :47-56. .

Shigenobu, S., Watanabe, H., Hattori, M., Sakaki, Y., and Ishikawa, H. (2000) Genome sequence of the endonuclear bacterial symbiont of aphids *Buchnera* sp. APS. Nature 407:886.

Sittenfeld, A., Uribe-Lorio, L., Mora, M., Nielsen, V., Arrieta, G., and Janzen, D. H. (2002) Does a polyphagous caterpillar have the same gut bacteria when feeding on different species of food plants? Rev. Biol. Trop. 50:547-560.

Sokal, R. R., and Rohlf, F. J. (1995) “Biometry” third edition, Freeman, New York.

Soria, C., and Mollema, C. (1995) Life-history components of western flower thrips on susceptible and resistant cucumber genotypes. Entomol. Exper. Appl. 74:177-184.

Spaulding, A. W., and Von Dohlen, C. D. (1998) Phylogenetic characterization and molecular evolution of bacterial endosymbionts in Psyllids (Hemiptera: Sternorrhychna). Mol. Biol. Evol. 15:1506-1513.

Stevenson, J. P. (1966) The normal bacterial flora of the alimentary canal of laboratory stocks of the desert locust, *Schistocerca gregaria* Forskål. J. Invertebr. Pathol. 8:205-211.

Stingl, U., Radek, R., Yang, H., and Brune, A. (2005) ‘*Endomicrobia*’: cytoplasmic symbionts of Termite gut protozoa form a separate phylum of Prokaryotes. Appl. Environ. Microbiol. 71:1473-1479.

Stouthamer, R., Breeuwer, J. A. J., Luck, R. F., and Werren, J. H. (1993) Molecular identification of micro-organisms associated with parthenogenesis. Nature 361:66-68.

Stouthamer, R., Breeuwer, J. A. J., and Hurst, G. D. D. (1999) *Wolbachia pipientis*: microbial manipulator of arthropod reproduction. Annu. Rev. Microbiol. 53:71-102.

Straif, S. C., Mbogo, C. N. M., Touré, A. M., Walker, E. D., and Kaufman, M. (1998) Midgut bacteria in *Anopheles gambiae* and *An. funestus* (Diptera: Culicidae) from Kenya and Mali. J. Med. Entomol. 35:222-226.

Subandiyah, S., Nikoh, N., Tsuyumu, S., Somowiyarjo, S., and Fukatsu, T. (2000) Complex endosymbiotic microbiota of the Citrus Psyllid *Diaphorina citri* (Homoptera: Psylloidae). Zool. Sci. 17:983-989.

Sutton, A. M., Becker, R. F., and Andaloro, J. T. (1983) Varietal resistance to onion thrips [Thysanoptera: Thripidae] in processing cabbage. J. Econ. Entomol. 76:85-86.

Swofford, D. (1990) PAUP 3.1.1. Champaign: Illinois Natural History Survey., Ill., USA.

Takahashi, K., Watanabe, K., and Sato, M. (1995) Survival and characteristics of ice nucleation-active bacteria on mulberry trees (*Morus* sp.) and in mulberry pyralid (*Glyphodes pyloalis*). Ann. Phytopathol. Soc. Jpn. 61:439-443.

Takatsuka, J., and Kunimi, Y. (2000) Intestinal bacteria affect growth of *Bacillus thuringiensis* in larvae of the oriental tea tortrix, *Homona magnanima* Diakonoff (Lepidoptera: Tortricidae). J. Invertrebr. Pathol. 76:222-226.

Tashiro, H., (1967) Selfwatering acrylic cages for confining insects and mites and detached leaves. J. Econ. Entomol. 60:354-356.

Terry, L. I. (1997) Host selection, communication, and reproductive behaviour. *In:* Thrips as Crop Pest (T. Lewis, ed.), p. 65-118. CAB Int., New York, N. Y., USA.

Terry, L. I., and Dyreson, E. (1996) Behavior of *Frankliniella occidentalis* (Thysanoptera: Thripidae) within aggregations, and morphometric correlates of fighting. Ann. Entomol. Soc. Am. 89:589-602.

Thao, M. L., Clark, M. A., Baumann, L., Brennan, E. B., Moran, N. A., and Baumann, P. (2000a) Secondary symbionts of Psyllids have been acquired multiple times. Curr. Microbiol. 41:300-304.

Thao, M. L., Moran, N. A., Abbot, P., Brennan, E. B., Burckhardt, D. H., and Baumann, P. (2000b) Cospeciation of Psyllids and their primary prokaryotic endosymbionts. Appl. Environ. Microbiol. 66:2898-2905.

Thao, M. L., Gullan, P., and Baumann, P. (2002) Secondary (g-*Proteobacteria*) endosymbionts infect the primary (ß-*Proteobacteria*) endosymbionts of the Mealybugs multiple times and coevolve with the host. Appl. Environ. Microbiol. 68:3190-3197.

Thao, M. L., and Baumann, P. (2004) Evidence for multiple acquisition of *Arsenophonus* by Whitefly species (Sternorrhychna: Aleyrodidae) Curr. Microbiol. 48:140-144.

Theunissen, J., and Legutowska, H. (1991) *Thrips tabaci* Lindeman (Thysanoptera, Thripidae) in leek: within plant distribution. J. Appl. Entomol. 112:309-316.

Thibout, E., Guillot, J. F., Ferary, S., Limouzin, P., and Auger, J. (1995) Origin and identification of bacteria which produce kairomones in the frass of *Acrolepiopsis assectella* (Lep., Hyponomeutidea). Experientia 51:1073-1075.

Thimm, Th., Hoffmann, A., Borkott, H., Much, J. Ch., and Tebbe, Chr. (1998) The gut of the soil microarthropod *Folsomia candida* (Collembola) is a frequently changeable but selective habitat and a vector for microorganisms. Appl. Environ. Microbiol. 64:2660-2669.

Thompson, J. N. (1987) Symbiont-induced speciation. Biol. J. Linnean Soc. 32:385-393.

Thompson, J. N. (1996) Trade-offs in larval performance on normal and novel hosts. Entomol. Exp. Appl. 80:133-139.

Thongaram, T., Hongoh, Y., Kosono, S., Ohkuma, M., Trakulnaleamsai, S., Noparatnaraporn, N., and Kudo, T. (2005) Comparison of bacterial communities in the alkaline gut segment among various species of higher termites. Extremophiles 9:229-238.

Toda, S., Murai, T., and Moreshita, M. (2007) Intra-specific genetic variation in *Thrips tabaci.* J. Insect Science May 4, 2007, Meeting Abstract. .

Treves, D. S., and Martin, M. M. (1994) Cellulose digestion in primitive hexapods: effect of ingested antibiotics on gut microbial populations and gut cellulase levels in the firebrat, *Thermobia domestica* (Zygentoma, Lepismatidae). J. Chem. Ecol. 20:2003-2020.

Trichilo, P. J., and Leigh, T. F. (1986) Predation on spider mite eggs by the western flower thrips, *Frankliniella occidentalis* (Thysanoptera: Thripidae), an opportunist in a cotton agroecosystem. Environ. Entomol. 15:821-825.

Tsiropoulos, G. J. (1976) Bacteria associated with the walnut husk fly, *Rhagoletis completa*. Environ. Entomol. 5:83-86.

Tsuchida, T., Koga, R., Meng, X. Y., Matsumoto, T., and Fukatsu, T. (2005) Characterization of a facultative endosymbiotic bacterium of the Pea Aphid *Acyrtosiphon pisum.* Microbiol. Ecol. 49:126-133.

Ullman, D. E., Westcott, D. M., Hunter, W. B., and Mau, R. F. L. (1989) Internal anatomy and morphology of *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) with special reference to interactions between thrips and tomato spotted wilt virus. Int. J. Insect Morphol. Embryol. 18:289-310.

Ullman, D. E., Cho, J. J., Mau, R. F. L., Westcott, D. M., and Custer, D. M. (1992) A midgut barrier to tomato spotted wilt virus acquisition by adult western flower thrips. Phytopathology 82:1333-1342.

Ullman, D. E., German, T. L., Sherwood, J. L., Westcott, D. M., and Cantone, F. A. (1993) *Tospovirus* replication in insect vector cells: Immunocytochemical evidence that the non-structural protein encoded by the S RNA of tomato spotted wilt tospovirus is present in thrips vector cells. Phytopathology 83:456-463.

Ullman, D. E., Sherwood, J. L., and German, T. L. (1997) Thrips as vectors of plant pathogens. *In:* Thrips as Crop Pest (T. Lewis, ed.), p. 539-566. CAB Int., New York, N. Y., USA.

Ulrich, R. G., Buhtala, D. A., and Klug, M. J. (1981) Microbiota associated with the gastrointestinal tract of the common house cricket, *Acheta domestica*. Appl. Environ. Microbiol. 41:246-254.

Vala Salvador, F. De Freitas, Van Opijnen, T., Breeuwer, J. A. J., and Sabelis, M. W. (2003) Genetic conflicts over sex ratio: mite-endosymbiont interactions. Am. Nat. 161:254-266.

Van Baalen, M., and Sabelis, M. W. (1995) The dynamics of multiple infection and the evolution of virulence. Am. Naturalist 146:881-910.

Van Baalen, M., and Janssen, V. (2001) Dangerous liaisons: the ecology of private interest and common good. Oikos 95:211-224.

Van Beneden, P. (1875) Les Commensaux et les Parasites. Bibliothèque Scientifique Internationale, Paris, France.

Van Borm, S., Buschinger, A., Boomsma, J. J., and Billen, J. (2002) *Tetraponera* ants have gut symbionts related to nitrogen-fixing rood-nodule bacteria. Proc. R. Soc. Lond. B 269:2023-2027.

Van de Heuvel, J. F. M. J., and Peters, D. (1989) Improved detection of potato leafroll virus in plant material and in aphids. Phytopathology 79:963-967.

Van de Wetering, F. (1999) Effects of Thrips Feeding on Tospovirus Transmission in Chrysanthemum. Ph D thesis, Wageningen Agricultural University, 120 pp. .

Van de Wetering, F., Goldbach, R. W., and Peters, D. (1996) Tomato Spotted Wilt Tospovirus ingestion by first instar larvae of *Frankliniella occidentalis* is a prerequisite for transmission. Phytopathology 86:900-905.

Van de Wetering, F., Van der Hoek, M. M., Goldbach, R. W., Mollema, C., and Peters, D. (1999) Variation in tospovirus transmission between populations of *Frankliniella occidentalis* (Thysanoptera: Thripidae). Bull. Entomol. Res. 89:579-588.

Van Dijken, F.R., Dik, M. T.A., Gebala, B., De Jong, J., and Mollema, C. (1994) Western flower thrips (Thysanoptera: Thripidae) effects on chrysanthemum cultivars: plant growth and leaf scarring in non-flowering plants. J. Econ. Entomol. 87:1312-1317.

Van Opijnen, T., and Breeuwer, J. A. J. 1999. High temperatures eliminate *Wolbachia*, a cytoplasmic incompatibility inducing endosymbiont, from the two-spotted spider mite. Exp. Appl. Acarol. 23:871-881.

Van Rijn, P. C. J., Mollema, C., and Steenhuis-Broers, G. M. (1995) Comparative life-history studies of *Frankliniella occidentalis* and *Thrips tabaci* (Thysanoptera: Thripidae) on cucumber. Bull. Entomol. Res. 85:285-297.

Van Rijn, P. C. J., Van Houten, Y. M., and Sabelis, M. W. (2002) How plants benefit from providing food to predators when it is also edible to herbivores. Ecology 83:2664-2679.

Via, S. (1990) Ecological genetics and host adaptation in herbivorous insects: the experimental study of evolution in natural and agricultural systems. Annu. Rev. Entomol. 35:421-446.

Vierbergen, G. (1995) International movement, detection, and quarantine of Thysanoptera pest. *In*: Thrips Biology and Pest Management (B. L. Parker, M. Skinner, and T. Lewis, eds.), p. 119-123. Plenum Press, New York, NY, USA.

Walsh, P. S., Metzger, D. A., and Higuchi, R. (1991) Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. Biotechniques 10:506-513.

Watanabe, K., Kawakita, H., and Sato, M. (1996) Epiphytic bacterium, *Erwinia ananas*, commonly isolated from rice plants and brown planthoppers (*Nlaparvata lugens*) in hopperbrun patches. Appl. Entomol. Zool. 31:459-462.

Watanabe, K., Katsuhara, M., Nakao, H., and Sato, M. (1997) Detection and molecular analysis of plant and insect-associated bacteria harboring aconitate isomerase involved in biosynthesis of trans-aconitic acid as antifeedant in brown planthoppers. Curr. Microbiol. 35:97-102.

Watanabe, K., Hara, W., and Sato, M. (1998) Evidence for growth of strains of the plant epiphytic bacterium *Erwinia herbicola* and transconjugation among the bacterial strains in guts of the silkworm *Bombyx mori.* J. Invertebr. Pathol. 72:104-111.

Watanabe, K., Abe, K., and Sato, M. (2000) Biological control of an insect pest by gut-colonizing *Enterobacter cloacae* transformed with ice-nucleation gene. J. Appl. Microbiol. 88:90-97.

Wells, M. L., Gitaitis, R. D., and Sanders, F. H. (2002) Association of Tobacco Thrips, *Frankliniella fusca* (Thysanoptera: Thripidae) with two species of bacteria of the genus *Pantoea*. Ann. Entomol. Soc. Am. 95:719-723.

Werren, J. H., Zhang, W., and Guo, L. R. (1995) Evolution and phylogeny of *Wolbachia*: reproductive parasites of arthropods. Proc. Roy. Soc. Lond. B 261:55-71.

Werren, J. H., and O'Neill, S. L. (1997) The evolution of heritable symbionts. *In*: Influential Passengers (S. L. O'Neill, A. A. Hoffmann, and J. H. Werren, eds.), p. 1-41. Oxford University Press, New York, NY, USA.

Whitehead, L. F., and Douglas, A. E. (1993) Populations of symbiotic bacteria in the parthenogenetic pea aphid (*Acyrtosiphon pisum*) symbiosis. Proc. Roy. Soc. Lond. B 254:29-32.

Whitfield, A. E., Ullman, D. E., and German, T. L. (2005) Tospovirus-Thrips interactions. Annu. Rev. Phytopathol. 43:459-489.

Wigglesworth, V. B. (1936) Symbiotic bacteria in a bloodsucking insect, *Rhodnius prolixus* Stal. Parasitology 28:284-289.

Wijkamp, I. (1995) Virus-Vector Relationships in the Transmission of Tospoviruses. Ph D thesis Wageningen Agricultural University, 143 pp. .

Wijkamp, I., and Peters, D. (1993) Determination of the median latent period of two tospoviruses in *Frankliniella occidentalis*, using a novel leaf disk assay. Phytopathology 83:986-991.

Wijkamp, I., Van Lent, J., Kormelink, R., Goldbach, R. W., and Peters, D. (1993) Multiplication of tomato spotted wilt virus in its insect vector, *Frankliniella occidentalis*. J. Gen. Virol. 74:341-349.

Wijkamp, I., Almarza, N., Goldbach, R. W., and Peters, D. (1995) Distinct levels of specificity in thrips transmission of tospoviruses. Phytopathology 85:1069-1074.

Yamamura, N. (1993) Vertical transmission and evolution of mutualism and parasitism. Theor. Popul. Biol. 44:95-109.

Yudin, L. S., Cho, J. J., and Mitchell, W. C. (1986) Host range of western flower thrips, *Frankliniella occidentalis* (Thysanoptera: Thripidae) with special reference to *Leucaena glauca*. Environ. Entomol. 15:1292-1295.

Yudin, L. S., Tabashnik, B. E., Cho, J. J., and Mitchell, W. C. (1988) Colonization of weeds and lettuce by thrips (Thysanoptera: Thripidae). Environmental Entomology 17:522-526.

Zawirska, I. (1976) Untersuchungen über zwei biologische Typen von *Thrips tabaci* Lind. (Thysanoptera, Thripidae) in der VR Polen. Archiv Phytopathol. Pflanzenschutz 12:411-422.

Zchori-Fein, E., and Brown, J. K. (2002) Diversity of Prokaryotes associated with *Bemisia tabaciI (*Gennadius) (Hemiptera: Aleyrodidae). Ann. Entomol. Soc. Am. 95:711-718.

Zhao, G., Liu, W., Brown, J. M., and Knowles, Ch. O. (1994) Insecticide resistance in field and laboratory strains of western flower thrips (Thysanoptera: Thripidae). J. Econ. Entomol. 88:1164-1170.

Zook, D. (1998) A new symbiosis language. Symbiosis News 1:1-3.

Zur Strassen, R. (1986) *Frankliniella occidentalis* (Pergande 1895), ein nordamerikanischer Fransenflügler (Thrysanoptera) als neuer Bewohner europäischer Gewächshäuser. Nachrichtenbl. Deut. Pflanzenschutzd. 38:86-88.

Zurek, L., Schal, C., and Watson, D. W. (2000) Diversity and contribution of the intestinal bacterial community to the development of *Musca domestica* (Diptera: Muscidae) larvae. J. Med. Entomol. 37:924-928.