

File ID uvapub:143711
Filename amc2006.60.pdf
Version unknown

SOURCE (OR PART OF THE FOLLOWING SOURCE):

Type article
Title Late onset axonal Charcot-Marie-Tooth phenotype caused by a novel myelin protein zero mutation
Author(s) H.M.E. Bienfait, C.G. Faber, F. Baas, A.A.W.M. Gabreëls-Festen, J.H.T.M. Koelman, J.E. Hoogendijk, J.J. Verschuuren, J.H.J. Wokke, M. de Visser
Faculty UvA: Universiteitsbibliotheek
Year 2006

FULL BIBLIOGRAPHIC DETAILS:

<http://hdl.handle.net/11245/1.426523>

Copyright

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content licence (like Creative Commons).



Late onset axonal Charcot-Marie-Tooth phenotype caused by a novel myelin protein zero mutation

H M E Bienfait, C G Faber, F Baas, A A W M Gabreëls-Festen, J H T M Koelman, J E Hoogendijk, J J Verschuuren, J H J Wokke and M de Visser

J. Neurol. Neurosurg. Psychiatry 2006;77;534-537
doi:10.1136/jnnp.2005.073437

Updated information and services can be found at:
<http://jnnp.bmj.com/cgi/content/full/77/4/534>

These include:

References

This article cites 9 articles, 5 of which can be accessed free at:
<http://jnnp.bmj.com/cgi/content/full/77/4/534#BIBL>

Rapid responses

You can respond to this article at:
<http://jnnp.bmj.com/cgi/eletter-submit/77/4/534>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[Genetics](#) (3952 articles)
[Neuromuscular disease](#) (353 articles)

Notes

To order reprints of this article go to:
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *Journal of Neurology, Neurosurgery, and Psychiatry* go to:
<http://www.bmjournals.com/subscriptions/>

SHORT REPORT

Late onset axonal Charcot-Marie-Tooth phenotype caused by a novel myelin protein zero mutation

H M E Bienfait, C G Faber, F Baas, A A W M Gabreëls-Festen, J H T M Koelman, J E Hoogendijk, J J Verschuuren, J H J Wokke, M de Visser

J Neurol Neurosurg Psychiatry 2006;**77**:534–537. doi: 10.1136/jnnp.2005.073437

A late onset axonal Charcot-Marie-Tooth phenotype is described, resulting from a novel mutation in the myelin protein zero (MPZ) gene. Comparative computer modelling of the three dimensional structure of the MPZ protein predicts that this mutation does not cause a significant structural change. The primary axonal disease process in these patients points to a function of MPZ in maintenance of the myelinated axons, apart from securing stability of the myelin layer.

Charcot-Marie-Tooth disease (CMT) is clinically and genetically heterogeneous. Patients are categorised in the demyelinating (CMT1) or the axonal form (CMT2). Patients with CMT1 have motor nerve conduction velocities (MNCV) of the median nerve below 38 m/s, whereas those with CMT2 have median nerve MNCV faster than 38 m/s. Mutations in two Schwann cell expressed myelin genes—the myelin protein zero gene (*MPZ*) and the Connexin 32 gene (*CX32*)—may show mainly axonal changes both electrophysiologically and pathologically.¹ A similar discrepancy between genotype and expected phenotype may occur with mutations in the neurofilament light gene (*NEFL*), a gene expressed in neurones, as *NEFL* mutations are often associated with nerve conduction velocities in the demyelinating range.

MPZ is the major protein component of peripheral nervous system myelin, composed of an extracellular, an intracellular, and a transmembrane domain. The extracellular domain is an immunoglobulin-like domain which plays an important role in compaction and maintenance of the peripheral myelin.² *MPZ* mutations are associated with CMT1, Dejerine-Sottas syndrome (DSS), congenital hypomyelination syndrome (CH), and CMT2.¹

We report a novel point mutation in the extracellular domain of the *MPZ* gene in two families with axonal CMT.

METHODS

Patients

In a cohort of 20 families with an axonal CMT phenotype, one family with a new *MPZ* mutation was identified. This mutation was present in several asymptomatic adult patients. Subsequently, the same mutation was found in a second CMT2 family. We obtained medical records of relatives with a CMT phenotype of the first family and undertook DNA analysis, which confirmed the *MPZ* mutation in the extended family.

Molecular genetic analysis

The genes for PMP22, *MPZ*, and *CX-32* were analysed for mutations by single strand confirmation polymorphism analysis (SSCP). The whole coding region of these genes was polymerase chain reaction (PCR) amplified from

genomic DNA and screened for altered mobility on SSCP gels. Gels with 0% and 10% glycerol were used for analysis. Fragments with aberrant migration were sequenced with BigDye terminator chemistry.

Sural nerve biopsy

Sural nerve biopsy specimens from two affected family members (one from each family) were available. Sural nerve biopsies were prepared for light and electron microscopic examination using standard techniques. The cluster ratio, defined as the number of clusters per 1000 myelinated fibres, was assessed by counting the number of clusters (three or more closely packed myelinated fibres) on the electron microscopic prints, divided by the number of myelinated fibres × 1000 in the same prints.

RESULTS

Patients


Clinical and electrophysiological data are listed in table 1. The pedigrees are shown in fig 1. Apart from H11.22 (father of H11.11), the Tyr82His mutation was confirmed in all patients. The phenotype is mild in most affected individuals and apart from a 34 year old female patient (H11.15) family members younger than 41 years were asymptomatic. Note that patient H11.11 did not have any complaints such as weakness, walking difficulty, or sensory symptoms and is therefore marked as asymptomatic, although clinical examination revealed minor weakness of toe flexor and extensor muscles and distal sensory abnormalities. Only one patient (H11.05) used a cane and boots for walking. Foot deformities, including pes cavus, pes planus, or claw toes, were found in six patients. Pes cavus was present in one asymptomatic patient only. There was no deafness co-segregating with the disease. Pupillary function was normal and lancinating pain was not mentioned.

The mean median nerve motor conduction velocity (MNCV) was 46 m/s and the mean ulnar nerve MNCV was 54 m/s. Median nerve MNCV was slower in symptomatic patients (mean 42 m/s) than in asymptomatic patients (mean 51 m/s) and was in the demyelinating range in two. Needle EMG of leg muscles showed spontaneous muscle fibre activity in three symptomatic patients, reinnervation signs in all but one patient (H11.08), and a reduced recruitment pattern in seven patients.

Molecular genetic analysis

An abnormal migration pattern was observed for the PCR fragment of exon 3 of the *MPZ* gene. The sequence of the fragment containing exon 3 on the *MPZ* gene was determined by dideoxy sequencing and a c.T244C substitution was found.

Abbreviations: CMT, Charcot-Marie-Tooth disease; DSS, Dejerine-Sottas syndrome; MNCV, motor nerve conduction velocity; SSCP, single strand confirmation polymorphism analysis



resulting in a p.Tyr82His substitution. This DNA sequence alteration was not found in over 200 unrelated control chromosomes tested.

Computer model prediction by SWISS-Model, based on the information of the crystal structure of MPZ protein, showed that the Tyr82His mutation does not cause a significant change in the tertiary structure of the MPZ protein.³

Sural nerve biopsy

Sural nerve biopsies were undertaken in patient H11.05 at the age of 46 years and in patient U01.02 at the age of 52. Both showed a marked loss of myelinated fibres (48% and 53%, respectively, of age matched control values), predominantly large diameter myelinated fibres (fibres >8 µm amounted to 15% and 4%, respectively, of age matched control values). Several clusters of regenerated myelinated axons were present. Cluster ratios were 29.7 and 50.2, respectively. No active axonal degeneration was observed, nor were signs of segmental demyelination, remyelination, or onion bulb formation. Electron microscopy revealed no structural abnormalities of myelin, Schwann cells, unmyelinated axons, or axon cytoskeleton.

DISCUSSION

Up to now, over 100 different mutations in the *MPZ* gene have been identified, from which approximately 10–15% show an axonal phenotype (Inherited Neuro-pathies Mutation Database: <http://www.molgen.ua.ac.be/CMTMutations/default.cfm>). How mutations in the *MPZ* gene cause mainly axonal injury is as yet not understood. We describe two families with a late onset axonal CMT phenotype caused by a Tyr82His mutation in the *MPZ* gene. Associated features such as deafness, lancinating pains, and pupillary abnormalities that were described in several axonal *MPZ* gene mutations were not present in our patients.^{4–5} Sensory symptoms seem to be more prominent in late onset axonal CMT caused by *MPZ* mutations.⁶ In our patients symptoms at onset were often sensory, but positive sensory features were not prominent. In all reports

regarding an axonal phenotype associated with *MPZ* mutations, disease onset is in or beyond the fourth decade, as in our families.^{1–6} Nearly all family members below age 40 in our study are in the presymptomatic stage. Electrophysiological examination revealed only minor abnormalities in these asymptomatic individuals and therefore cannot reliably be used for case ascertainment in this stage of the disease. The rare occurrence of pes cavus is in line with an actual disease onset beyond the first years of life. The mild phenotype, together with the lack of pes cavus in most patients with this mutation, may cause confusion with acquired neuropathy if the hereditary nature of the disease is not recognised.

The Tyr82His mutation in the *MPZ* gene has not been reported previously. However, mutations of tyrosine 82 have been described in CMT1 and DSS patients with onset in (early) childhood and very slow MNCVs.^{7–9} In these cases, the tyrosine 82 was substituted for by cysteine. The difference in phenotype can be attributed to the nature of the mutation. The replacement of tyrosine by cysteine in the extracellular part of the protein will have a large impact on the tertiary structure and thus on myelin structure and function owing to the introduction of novel disulphide bridges. In case of a tyrosine to histidine mutation, modelling studies predicted that our Tyr→His mutation does not cause a significant structural change, which is in line with the lack of primary demyelinating features in our patients. Instead, nerve conduction studies were compatible with a diagnosis of CMT2 in most patients. The electrophysiological findings in our patients and the absence of a primary demyelinating process in two nerve biopsies indicate that axonal degeneration and regeneration is the primary pathophysiological basis of the CMT phenotype caused by the Tyr82His mutation. This suggests a function of MPZ in the maintenance of myelinated axons, apart from securing stability of the myelin layer.

ACKNOWLEDGEMENTS

We thank J P Vreijling for DNA sequence analysis and J Bekker for the pedigree drawings. This study was supported by funds from the Prinses Beatrix Fonds, The Hague.

.....
Authors' affiliations

H M E Bienfait, J H T M Koelman, M de Visser, Department of Neurology, Academic Medical Centre, University of Amsterdam, Amsterdam, Netherlands

J E Hoogendijk, J H J Wokke, University Medical Centre Utrecht, Utrecht, Netherlands

C G Faber, University Hospital Maastricht, Maastricht, Netherlands

J J Verschuuren, Leiden University Medical Centre, Leiden, Netherlands

A A W M Gabreëls-Festen, Institute of Neurology, University Medical Centre St Radboud, Nijmegen, Netherlands

F Baas, Neurogenetics Laboratory, Academic Medical Centre, University of Amsterdam, Netherlands

Competing interests: none declared

Correspondence to: Professor M de Visser, Department of Neurology H2-222, Academic Medical Centre, PO Box 22660, 1100 DD Amsterdam, Netherlands; M.deVisser@amc.uva.nl

Received 4 June 2005

In revised form 9 December 2005

Accepted 9 December 2005

REFERENCES

- 1 **Shy ME**, Jani A, Krajewski K, *et al*. Phenotypic clustering in MPZ mutations. *Brain* 2004;**127**:371–84.
- 2 **Lemke G**. The molecular genetics of myelination: an update. *Glia* 1993;**7**:263–71.
- 3 **Schwede T**, Kopp J, Guex N, *et al*. SWISS-MODEL: an automated protein homology-modeling server. *Nucleic Acids Res* 2003;**31**:3381–5.
- 4 **Chapon F**, Latour P, Diraison P, *et al*. Axonal phenotype of Charcot-Marie-Tooth disease associated with a mutation in the myelin protein zero gene. *J Neurol Neurosurg Psychiatry* 1999;**66**:779–82.
- 5 **De Jonghe P**, Timmerman V, Ceuterick C, *et al*. The Thr124Met mutation in the peripheral myelin protein zero (MPZ) gene is associated with a clinically distinct Charcot-Marie-Tooth phenotype. *Brain* 1999;**122**:281–90.
- 6 **Auer-Grumbach M**, Strasser-Fuchs S, Robl T, *et al*. Late onset Charcot-Marie-Tooth 2 syndrome caused by two novel mutations in the MPZ gene. *Neurology* 2003;**61**:1435–7.
- 7 **Himoro M**, Yoshikawa H, Matsui T, *et al*. New mutation of the myelin P0 gene in a pedigree of Charcot-Marie-Tooth neuropathy 1. *Biochem Mol Biol Int* 1993;**31**:169–73.
- 8 **Silander K**, Meretoja P, Juvonen V, *et al*. Spectrum of mutations in Finnish patients with Charcot-Marie-Tooth disease and related neuropathies. *Hum Mutat* 1998;**12**:59–68.
- 9 **Boerkoel CF**, Takashima H, Garcia CA, *et al*. Charcot-Marie-Tooth disease and related neuropathies: mutation distribution and genotype-phenotype correlation. *Ann Neurol* 2002;**51**:190–201.