Downloaded from UvA-DARE, the institutional repository of the University of Amsterdam (UvA) http://hdl.handle.net/11245/2.30186

File ID uvapub:30186 Filename HS150805 Version unknown

SOURCE (OR PART OF THE FOLLOWING SOURCE):

Type book chapter

Title Non parametric population classification

Author(s) A.W.M. Smeulders

Faculty UvA: Universiteitsbibliotheek

Year 1986

FULL BIBLIOGRAPHIC DETAILS:

http://hdl.handle.net/11245/1.423961

Copyright

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content licence (like Creative Commons).



A078355265 (16)NCC/IBL AANVRAAGBON KOPIE MONOGRAFIE EGB 18-04-2005 Datum indienen : 17-04-2005 10:33 Clearing House 5493-1 Datum plaatsen : 17-04-2005 10:33 Aanvrager : 0004/9998 Aanvraagident : Aanvragerident : 0004/9999 Eindgebruiker : 041631433 Telefoonnummer: 050-3635057 Cooperatiecode : R : Elektronisch Leverwijze Fax Ftp E-Mail : m.s.van.delden@rug.nl Ariel Plaatscode : 221777210 ; MB 133 e023 ; ; Origineel gestuurd Niet beschikbaar Copie gestuurd Uitgeleend Overige (8) [] Wordt niet uitgeleend Bibliografisch onjuist Nog niet aanwezig Niet aanwezig Bij de binder (5) Verzamelnota volgt. A078355265 18-04-2005 NCC/IBL AANVRAAGBON KOPIE MONOGRAFIE EGB Datum indienen : 17-04-2005 10:33 9560-1 UvA Keur Aantal Datum plaatsen: 17-04-2005 10:33 UB Groningen Broerstraat 4 : 0004/9998 Aanvrager 9700 AN Groningen Aanvraagident : Aanvragerident: 0004/9999 Eindgebruiker : 041631433 tav PPN Titel : 221777210 : FIRST QUINQUENNIAL REVIEW 1981-1986 Titel : Backer, E. (ed.) Auteur Deel/Supplem. : Dutch Society for Pattern Recognition and Image Processing Corporatie Jaar/Editie : 1986 Extern nummer : Uitgave : Pijnacker DEB Serie/Sectie Pag-ISSN/ISBN : 90-6231-144-X : 221777210 ; MB 133 e023 ; ; Plaatscode Jaar Volume Aflevering Aanvraagident. Eindgebruiker : 041631433 Aanvragerident. : UVA KEUR (UB GRONINGEN)

Bron Opmerking 136060 : arno ID:

: 39-53

Auteur

Artikel

Bladzijden

A.W.M. SMEULDERS

Department of Pathology, and the Department of Medical Informatics, Free University, De Boelelaan 1117, 1081 HV Amsterdam, The Netherlands


SUMMARY

Population classification is a set of statistical techniques to classify populations on basis of observations of their constituing members. This decision making involves a cascade of classifiers: one for objects, and, based on the resulting classifications, one for populations. Also a rule is to be given how many objects should be analysed before the population is decidable. In search for a more integral view of the problem the population function is introduced. Consideration of the population function implies that rather then fixing the object classifier a priori, it is more efficient and more accurate to extend the classifier to a distinct range of the feature vector. The concept of sequential classification is defined as well. Population classification performance is favourably compared to /1,2,3,5,6/.

1. INTRODUCTION

Population classification is the classification of a population on basis of observations on its members. Population classification deviates from usual decision making procedures in the two step classification procedure that is employed. In the first step, members of a population each are classified into one of the member classes. After the analysis of a certain number of members, in the second step the population itself is classified into one of the population classes. Population classification is a statistical technique with applications in many different

fields such as industrial inspection, quality control and quality assessment of lots. For conceptual convenience, we will adopt here the terminology from the cytology field, where a specimen may be conceived as a population of cells.



In figure 1, the two step classification procedure is depicted. It is crucial in this cascade of classifiers that the accumulated outcome of the cell classification is the input of the specimen classification. So, an alteration in the cell classifier may influence the specimen classification result. In the analysis of specimens, the prime interest is not in the performance of the cell classifier on its own, but in its performance in respect to the specimen classification.

Let the cell of the I-th specimen be numbered by index i with $i=1,\ldots,n_{\mathrm{I}}$. A feature vector w_i is computed per cell, giving a list of n_{I} observations. The cascade of classifiers is composed of three tunable entities:

- 1. The cell classifier (\bar{w}_t) provides a set of rules to classify a cell characterized by \bar{w}_i . The cell is classified into one or more cell classes k with k=1,...,m. The cell classifications are accumulated in the cell class occupancy vector f. It should be noted, that in the results
- use a cell classifier. 2. The number rule (n_t) controls the minimum number of cells to be classified before classifying the specimen.

of this paper will suggest that it may be profitably not to

3. The specimen classifier (f_t) provides a set of rules to classify a specimen based on f after the analysis of n_t cells. The specimen is classified in specimen class K with $K=1,\ldots,M$.

The objective of statistical population analysis is to design a procedure such that the specimen classification error is acceptable and the number of cells to classify is minimized by controlling these three entities. It is important to note that a specimen may either be erroneously classified by the fact that only a limited number of cells are analysed or by the fact that the occupancy vector \mathbf{f}_t of a specimen, when compared to \mathbf{f} , leads to erroneous classification. The first error, the sample error, is related to the system efficiency and may be avoided by analysing more cells. The second error is related to accuracy of the system and is intrinsic to the nature of the problem. For a fixed $\overline{\mathbf{w}}_t$ and \mathbf{f}_t , it cannot be cured by analysing more cells.

Castleman and White were the first to design a procedure for cascaded classification. In a series of a papers, using a simplified model, they have given guidelines how to address the cell classifier /l/, how to address the specimen classifier /2/, and how the proportionality of different classes of cells influences the performance of the specimen classifier /3/, once the cell classifier has been fixed. All references are restricted to M=2 with K=l: 'normal specimens' and K=2: 'abnormal specimens'. In /l/ and /3/, the number of cell classes is restricted to m=2. In that case, we will call cells classified into class k=1 'non-events', and cells into class k=2, 'events'. In /2/ there is no restriction to the number of cell classes.

The procedure of Castleman and White is discussed here, without loss of generality, for m=2. They consider a previously settled cell classifier such that a fraction (ϵ_1) of normal cells (k=1) is erroneously classified into class k=2, and fraction (ϵ_2) of abnormal cells erroneously into k=1. Secondly, the presumption is made that specimens of the normal specimen class (K=1) entirely consists of k=1-cells. Also, it is presumed that abnormal specimens (K=2) all have the same, fixed, fraction p of k=2-cells, and a fraction 1-p of k=1-cells. Let q_p for K=

1,2 be the proportion of events from a specimen of class K, then we have $q_1 = \varepsilon_1$ and $q_2 = \varepsilon_1 + p(1-\varepsilon_1-\varepsilon_2)$. Both q_1 and q_2 are thus constants for all specimens of class K.

It should be noted that the discrimination of ϵ_1 and ϵ_2 from p is not necessary. The same result would have been obtained by the presumption that the cell classifier produces a proportion of q_1 events for normal specimens and q_2 events for abnormal ones.

Under these presumptions, the classification procedure reduces to the recognition whether an observed proportion \hat{q} of an unknown specimen originates from a specimen with q_1 or q_2 events. As \hat{q} is based on a cell, \hat{q} is distributed binomially around q_K . The binomial distribution is, with a very good approximation, replaced by a Gaussian distribution with mean q_K and standard deviation $s_K = \sqrt{q_K(1-q_K)/n}$. Let $f_e(x)$ be the error function

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{q}^{\infty} e^{-t^2} dt$$
 (1a)

and $E_K^{}$, K=1,2, be the tolerated fractions of specimens classified wrong. Define A and B implicitly by:

$$E = f(B)$$
 and $E = f(A)$. (1b)

The minimum number of cells to be analysed (n_t) and the specimen classification threshold (q_t) may now be solved as follows:

$$n_{t} = \left(\frac{A\sqrt{q_{2}(1-q_{2})} + B\sqrt{q_{1}(1-q_{1})}}{q_{2}-q_{1}}\right)^{2}$$
 (2)

$$q_{t} = \frac{A q_{2}\sqrt{q_{1}(1-q_{1})} + B q_{1}\sqrt{q_{2}(1-q_{2})}}{A\sqrt{q_{1}(1-q_{1})} - B\sqrt{q_{2}(1-q_{2})}}$$
(3)

In summary, the proposed procedure is:

- l. Select cell features \bar{w} and a cell classifier \bar{w}_{t} , see [1].
- ?. Select a value for \mathbf{q}_1 representative for normal specimens on basis of experience with this cell classifier. Similarly, choose a \mathbf{q}_2 . Compute \mathbf{n}_t and \mathbf{q}_t .

3. In the operational phase, analyse n_t cells for each specimen and find \hat{q} . If $\hat{q} < q_t$ then decide "normal specimen" else $\hat{q} \ge q_t$ decide "abnormal specimen".

Although the procedure is attractive in the sense that it provided a relationship between the cell clasifier and the specimen classifier, the presumptions tend to oversimplify reality. As a consequence all information in $\bar{\mathbf{w}}_i$ is reduced two parts above and below $\bar{\mathbf{w}}_t$. All other information in $\bar{\mathbf{w}}$ is lost. Secondly, in many practical cases, it is in conflict with reality to presume that a class of specimens can be represented by a fixed proportion of events. Thirdly, it is not necessary to select n a priori. Wald /4/ has demonstrated that \mathbf{n}_t is most efficiently computed during the analysis of the specimen, as was also recognised in /2/. Finally, chosing a cell classifier prior to and separately from the establishment of the specimen classifier is not optimal, as will be demonstrated later on.

An alternative approach was followed by Tanaka /5/, acquiring the values of one cell feature in a cumulative histogram q(v). The procedure that was used, was as follows:

- 1. One cell feature, v, is selected.
- 2. In the learning phase, derive histograms $q_{K}\left(v\right)$ representing the K-th specimen class.
- 3. In the operational phase, establish after the analysis of n cells, the specimen histogram $\hat{q}_K(v)$, and apply the non-parametric Kolmogorov Smirnov test for the difference of two histograms.

For K=1,2:

$$v_t = v \text{ where max } | \hat{q}_n(v) - q_K(v) |$$
 (4a)

$$n_{t} = \frac{(f_{KS}(E_{K}^{S}))^{2}}{(\hat{q}_{n}(v_{t}) - q_{K}(v_{t}))^{2}}$$
(4b)

if
$$n > n_{t}$$
 assign I to class K (4c)

else assign I not to a class.

In this decision rule $f_{KS}(E_K^S)$ is a factor based on the confidence level E_K of the Kolmogorov Smirnov test. The values of f_{KS} may be found in standard statistical tables /11/.

The analysis with a non-parametric test, only permits the

use of a one dimensional cell feature. Non-parametric tests on the difference of two distributions, do not exist for more than one dimension, as the more dimensional space cannot be uniquely ordered.

More dimensional cell features (\bar{w}) can be used still, by using non-linear mapping (M) of the cell features onto what is called the axis of cell atypia. The index (v)

$$\mathbf{v} = \mathbf{M}(\mathbf{w}) \tag{5}$$

winds through the feature space, e.g. according to /6/ or /7/. In reference /6/, the axis is a result of a curve fitting procedure through a number of ordered cell classes represented by their covariance matrices, derived from a learning phase. The ordering of the cell classes permits the definition of a scale, called index of atypia, on the axis. De facto, this axis can be regarded as a way to order the cell feature space and classes, giving a meaning to the feature values in the context of the classification problem. The index of atypicality is an ordinal scale, and can on its own be regarded as a one dimensional cell feature. In /6/, no hint is given how to use the index in a specimen classification procedure.

A conceptually different way to reduce the more dimensional cell feature space into a one dimensional cell scale, is making use of a posteriori probability of a cell classifier. Let M now denote a (set of) cell classifier(s), and let v now denote the resulting a posteriori probability of a cell to belong to cell class k=1, then v also is specified by equation 5. The vector v is an ordinal scale cell feature. For some problems, for M the Mahalanobis distance to the class of normal cells may be taken as a measure for the deviation of the cell from the normal /9/. In this concept and the previous one, a cell feature v is the residue from the more dimensional feature $\bar{\mathbf{w}}$ by an operation M.

Ott /8/ also applies a mapping from the more dimensional cell space onto a triangulated cell class space. Specimen features are defined on this cell class space as input to the specimen classifier. No hints are, however, given how to

generally define specimen features or how to relate the number of cells to be analysed on a specimen to the classification error.

In the present paper, it is pointed out how the distribution of v relates to the specimen classification error by the introduction of the population function, see also /10/.

2. THE POPULATION FUNCTION

Consider an arbitrary specimen I of specimen class K. For a one dimensional cell feature v, which may be the result of a mapping (v=M(w)), call the probability density function p(v). A convenient form to consider is the cumulative density function

$$q(v) = \int_{v}^{\infty} p(x) dx.$$
 (6)

It is interesting to relate q(v) to a cell classifier v^{*} of the following type, see figure 2.

In this context, q(v') denotes the fraction of cells that will be classified into class k=2 (events), when v' would have been used as the cell classifier. The fraction non-events accordingly is 1-q(v'). When v' is varied over the feature scale, q(v) denotes the performance for any cell classifier v on the given specimen.

For one cell classifier v', the fraction of events may vary from specimen to specimen, even within one class of specimen. This fraction q(v') thus is a variable itself, for which within one class of specimens K, a pdf may be formed: $P_K(q(v'))$. This function indicates the probability to observe a fraction q(v') of events when analysing a specimen of class K with cell classifier v'. Obviously the shape of $P_K(q(v'))$ is completely dependent on the nature of problem and is independent of q(v). Variation of v' over the feature scale yields the result for any feature: $P_K(v,q(v))$. This is the population function, giving a




Figure 2: The proportion of events (cells classified into k=2) resulting from a cell classifier v' indicated by shading in the graph of p(v) and the cumulative function q(v).

relationship between v and q(v) for population K. The integral form of P_{K} over q:

$$Q(v,q(v)) = \int_{0}^{q} P(v,x(v))dx$$
(8)

relates an arbitrary cell classifier to the obtainable specimen classification error as follows. Consider a specimen classifier q' such that:

if
$$q_{\rm T}\!\!<\!\!q'$$
 assign specimen I to specimen class K=1, (9) else $q_{\rm T}\!\!\ge\!\!q'$ assign specimen I to class K=2.

For a given cell classifier v' and a given specimen classifier $q',\ Q_K(v',q'(v'))$ denotes the probability of a specimen to be classified into class K=1. The function $1-Q_1(v,q(v))$ thus is the specimen classification error function for K=1-specimens, and the function $Q_2(v,q(v))$ is the specimen error-function for K=2-specimens.

The population function can be used in specimen classification procedures by considering for specimen class K=1,2, the population functions $P_K(v,q(v))$ as illustrated in figure 3a,b. For each value v' of v, the Bayes-rule implies that the specimen classifier with smallest total error is $q_B(v')$ at the point where $P_1(v',q(v')) = P_2(v',q(v'))$. Varying v' again

feature scale gives a curve $q_B(v)$, indicating for each sociated best specimen classifier. The cumulative n function is displayed in figure 3b. For each v', w is positioned at the location of maximum vertical between $Q_1(v',q(v'))$ and $Q_2(v',q(v'))$. In figure 3c, section of $q_B(v)$ with $Q_K(v,q(v))$ is indicated. The specimen classification error, is indicated by (v) and $Q_1(v,q(v))$. In the example of figure 3c, for er part of the feature vector v the population v is alishable from population 2, and consequently on that error functions have a value .5.

al conclusions may be drawn from figure 3c. Firstly, be a part of the feature scale v where no or little ation is possible between class-K=1 and class-K=2. Secondly, for the part of the scale where ion is possible, in figure 3c denoted by the region especimen classifiction error is $1-Q_1(v,q(v))$ for K=1 q(v)) for K=2. This error is due to the nature of the expressed in the shapes of $Q_K(v,q(v))$ and can for ture not be avoided. Thirdly, if desired, the single classifier v_0 can be selected by taking

$$\max_{\mathbf{v}, \mathbf{q}} | Q_1(\mathbf{v}, \mathbf{q}_B(\mathbf{v})) - Q_2(\mathbf{v}, \mathbf{q}_B(\mathbf{v})) |.$$
 (10)

L see lateron, it may still be advantageous not to zell classifier a priori.

NG PHASE

a far, we have considered cell classifiers and specimen is for the true population functions. In principle, we ispose of the true $P_K(v,q(v))$. Instead we have to rely ions or simplifications, based on a limited series of ins $\hat{q}_I(v)$, $I=1,\ldots,N_K$ for K=1,2. These observations the learning set of specimens.

> rous simplification of the population function as used has already been discussed in the introduction. The two independent simplifications. First, one cell

classifier is selected, reducing $P_K(v,q(v))$ to a one dimensional function. Secondly, as q(v) is assumed to be equal for all specimens of the same class, $P_K(v,q(v))$ then further reduces to a delta-pulse. Especially the second simplification is in conflict with common classification problems. Note, that in the context of this model no overlap occurs between the approximations for the population functions. As a consequence, there is no preference for a specimen classifier qt on basis of the assumed properties of the two populations at hand other than $q_1 < q_2$. The expression for q_t given in eq.(3) is derived on basis of the finite number of cells only; a subject we will come to discuss in the next section.




Figure 3a,b: The population function $P_K(v,q(v))$ shown left for one specimen class. For the ease of display P(v,q(v)) has been chosen here unimodal, Gaussian shaped for all cross-sections over v. Note that, in general, not all cross-sections of P(v,q(v)) are necessarily Gaussian shaped or even unimodal, nor have a similar shape. On the right side a pair of joint probability functions $P_K(v,q(v))$ is given. In the cross-section at the front, the Bayes specimen classifier for that cross-section is indicated by $q_R(v)$.

Another, much better estimation, $\hat{P}_K(v,q(v))$ for the population function is the following Parzen-estimation procedure /10/:

$$P_{K}(v,q(v)) = \frac{1}{N_{K}} \sum_{I \in K} \frac{1}{S_{I} \sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{q(v)-q}{S_{I}} \frac{(v)}{S}\right)^{2}}$$
 with (11)

$$S_{I} = \sqrt{\frac{q_{I}(v)(1-q_{I}(v))}{N_{I}}}$$
 (12)

Having established $P_K(v,q(v))$ at the end of the learning phase, $q_t(v)$ can be determined as an estimate for $q_B(v)$. $q_t(v)$ is

implicitly defined by $\hat{P}_1(v,q_t(v))=\hat{P}_2(v,q_t(v))$. Analogously to the cumulative population function, Q(v,q(v)) gives error estimates for this specimen classifier. If E_K^B is the tolerated specimen classifier error fraction for class K due to overlap of $P_1(v,q(v))$ and $P_2(v,q(v))$, then a region v_{test} of v may be selected as follows:

$$v_{test} = v \text{ where } 1 - Q_1(v, q_t(v)) < E_1^B \text{ and. } Q_2(v, q_t(v)) < E_2^B$$
 (13)

If we could dispose of the cumulative probability density function $q_I(v)$ of an unknown specimen I, we could classify the specimen by comparing $q_t(v)$ to $q_I(v)$ for the region v_{test} . In that case, the error may be expected to be bounded by E_K^B . In reality, this specimen classification error is not the only source of error as shall be discussed in the next section.

4. SEQUENTIAL CLASSIFICATION

In the analysis of an unknown specimen I, a finite number n of cells is analysed, resulting in a cumulative histogram of feature values $\hat{q}_{-}(v)$. The histogram is a sample of an unknown




Figure 3c,d: The cumulative population functions $Q_K(v,q(v))$ on the left. The specimen classifier error as the intersection of $Q_K(v,q(v))$ with $q_R(v)$ on the right.

distribution $\hat{q}_1(v)$. A specimen is said to be undecidable when $q_n(v)$ may be a sample of $q_t(v)$ over the region v_{test} . In order to test the decidability a null hypothesis is formed stating that $\hat{q}_n(v)$ is a sample of $q_t(v)$. A hypothesis test on the difference of the two distributions is used. Two non-parametric tests are considered here.

The first to consider is the Kolmogorov Smirnov test. Let \mathbb{E}_K^S be the tolerated specimen fraction to be classified wrong due to finite cell sample size. This factor then serves as the level of confidence in the test, see eq.(4). The specimen decision rule based on the KS-test is:

$$v_t = v$$
 where max $|\hat{q}_n(v) - q_t(v)|$ (14a)
 v_{test}

$$n_{t} = \frac{(f_{KS}(E_{K}^{S}))^{2}}{(\hat{q}_{n}(v_{t}) - q_{t}(v_{t}))^{2}}$$
(14b)

if $\hat{q}_n(v) < q_t(v_t)$.and. $n > n_t$ assign I to class K=1

if
$$\hat{q}(v)\geq q(v)$$
 and $n>n$ assign I to class $K=2$ (14c)

else specimen I is undecidable.

Note the difference between eq.(4) and eq.(14). The null hypothesis used in /5/ was that $\hat{q}_n(v)$ originated from $q_1(v)$ or $q_2(v)$ which are example histograms of specimen class K=1 and K=2. In many problems, a class of specimens cannot reliably be represented by one example histogram. Doing so may lead to large classification errors. Alternatively, in eq. (14), $\hat{q}_n(v)$ is tested against $q_t(v)$, the critical function. When $\hat{q}_n(v)$ deviates significantly from $q_t(v)$, a decision can be taken with a known, bounded error probability.

A drawback of the KS-test is its aspecificity. Many samples are needed to demonstrate a significant difference. As a solution, the following test is proposed.

Consider one specimen class v'. Let the tolerated fraction of specimen to be classified wrong again be \mathbf{E}_K^S . As \mathbf{q}_t (v') is a fraction, samples of \mathbf{q}_t (v') based on n observations (cells) are distributed binomially with mean \mathbf{q}_t (v') and standard deviation similar to eq.(12). With good approximation, the binomial distribution may be replaced by the Gaussian one with same mean and standard deviation. A relation now may be found between the value of $\hat{\mathbf{q}}_n$ (v') needed to reject the hypothesis given \mathbf{q}_t (v'), n and \mathbf{f}_e (\mathbf{E}_K^S), see eq.(15).

The same test holds, varying v'over v. In that case $\textbf{q}_{1}(\textbf{v})$ is said to be decidable if one value v' is found for which

 $\hat{q}_{n}(v')$ significantly differs from $q_{t}(v')$.

The proposed test becomes:

For K=1,2

$$v_t = v \text{ where } \min_{t \in V_{test}} \{n_t(v)\}, \text{ with}$$
 (15a)

$$n_{t}(v) = \frac{q_{t}(v) (1-q_{t}(v)) (f_{e}(E_{K}^{S}))^{2}}{(q_{t}(v) - \hat{q}_{n}(v))^{2}}$$
(15b)

if $n > n_t(v_t)$ assign specimen I to class K=1 if $\hat{q}_n(v_t) < q_t(v_t)$ if $n > n_t(v_t)$ assign specimen I to class K=2 if $\hat{q}_n(v_t) > q_t(v_t)$ else $n \le n_t(v_t)$ and specimen I is undecidable.

The comparison of eq.(4) with eq.(14), can also be made for this test, comparing eq.(2) and (3) with eq.(15). Again, rather than comparing $\hat{q}_n(v)$ with one particular value $q_1(v)$ or $q_2(v)$ as was done in /1-3/, $\hat{q}_n(v)$ here is compared with $q_n(v)$. In practice, it is likely to lead to smaller classification errors.

It should be noted that the above tests could be used to compute a value for n to analyze on all unknown specimens. It is more profitable, however, to apply eq.(14) or eq.(15) sequentially. In that case, the entire population classification becomes:

- l. Define an operation M reducing the more dimensional cell feature space \bar{w} to a one dimensional vector /6,9/.
- 2. In the learning phase, estimate the population functions by eq.(11) and compute $q_t(v)$. Specify the tolerated error fractions due to specimen overlap and find v_{test} , eq.(13).
- 3. In the test phase, specify the tolerated error fractions due to the finite cell sample size. Update $\hat{q}_n(v)$ after the analysis of each cell and apply the following decision rule:

Evaluate eq.(14) or eq.(15)

If the specimen is decidable, stop the analysis, else proceed with the next cell.

It has been made pausible /10/, that the procedure based on eq.(15) is both more efficient, in the sense that it needs fewer cells to arrive at a conclusion, and more accurate in the sense that it makes fewer errors in the classification, than the

method of eq.(14). Also, this procedure is likely to be more efficient and accurate than the method based on eq.(2) and (3) /1-3/, and more accurate than the method of eq.(4) /5/. Model studies are currently being done to investigate these conjectures /12/.

5. CONCLUSION

In the foregoing, the population function $P_K(v,q(v))$ was introduced as a theoretical framework to describe the performance of cascade classifiers. Three items are important here, the cell (primary) classifier, the specimen (secondary) classifier and a rule prescribing how many cells need to be analysed before a specimen is decidable. The performance of a cascade classifier is established by its accuracy (specimen classification error) and its efficiency (number of cells to be analysed).

The population function is the equivalent of the probability density function for one-classifier problems. The equivalent of the Bayes-rule, optimizing overall accuracy, is for population classification given by eq.(10). Equation (10) will give the theoretical most accurate cell classifier selected prior to the analysis and the associated specimen classifier. This gives a theoretical solution to the problem raised in /1/ and /2/. From eq.(10) it is also clear that the cell and specimen classifier are best optimized in combination, not separately, as was done in /1,2/. A priori selected cell classifiers will be further discussed in /13/.

In practice, eq.(10) can only be approximated by a learning phase for which we propose the computation of the Parzen-estimation given in eq.(11), and the selection of a part of the cell feature as specified in eq.(13). The selection of this part of the feature vector implies that the specimen classification error is restricted to a bound, derived in /12/.

In spite of eq.(10), it is not necessary (even unprofitable) to select a cell classifier prior to the analysis, as was done in /1,2,3/. Bounding the tolerated specimen classification error to a specified fraction as proposed in eq.(13), fewer cells need

- - ---- - -

to be analysed when using sequential classification techniques. The technique differs from previous ones in that the difference is demonstrated with the threshold function q (v) between the two populations rather then the concordance with the examplatory population functions /3,4,5/. This is expected to lead to a higher accuracy. Presently, model studies are carried out to investigate the accuracy and efficiency of the proposed models in comparison to the existing ones /12/.

ACKNOWLEDGEMENT

G. Bosveld and R.P.W. Duin are gratefully acknowledged for critically reviewing the paper. The manuscript was carefully prepared for publication by D.I.M. de Jong.

REFERENCES

- 1. Castleman, K.R. and White, B.S., The tradeoff of cell classifier error rates, Cytometry 1 (1980) 601-613.
- 2. Castleman K.R. and White, B.S., Optimizing cervical specimen classifiers, IEEE trans. PAMI 2 (1980) 451-457.
- 3. Castleman, K.R. and White, B.S., The effect of abnormal cell proportion on specimen classifier performance, Cytometry 2 (1981) 155-158.
- 4. Wald, A., Sequential analysis, Dover Publications, New York (1947).
- 5. Tanaka, N., Ikeda, H., Ueno, T., Mukawa, A., Kamitsuma, K., Field test and experimental use of CYBEST model 2 for practical gynecologic mass screening, J. Anal. Quant. Cytol. 1 (1979) 122-126.
- 6. Bartels, P., Bibbo, M., Richards, D.L., Sychra, J.J., Wied, G.L., Patient classification based on cytologic sample profiles. I Basic measures for profile constructions, Acta Cytol. 22 (1978) 253-260.
- 7. Gelsema, E.S., Hunink, M., Queiros, C.E., Timmers, T., The use of correspondence analysis in the assessment of morphological changes during carcinogenesis, Cytometry 5 (1984) 463-468.
- 8. Ott, R., Schurman, J., Reinhardt, E.R., Bloss, W.H., Automated classification of cytological specimens based on features extracted from nuclei images, Pattern Recognition 13 (1981) 83-87.
- 9. Mayall, B. and Burger, G., Personal communications.
- 10. Smeulders, A.W.M., Pattern analysis of cervical specimens, Ph.D.-thesis, University of Leyden, The Netherlands (1983).
- 11. Beyer, W.H. et al, Handbook of probability and statistics, The Chemical Rubber co. Cleveland (1968).
- 12. Bosveld, G., Smeulders, A.W.M., Duin, R.P.W., In preparation.
- 13. Smeulders, A.W.M., Bosveld, G., Duin, R.P.W., Optimisation of cacade classifiers by means of the population function. In preparation.