Downloaded from UvA-DARE, the institutional repository of the University of Amsterdam (UvA)
http://hdl.handle.net/11245/2.44300

File ID	uvapub:44300
Filename	Contents
Version	unknown

SOURCE (OR PART OF THE FOLLOWING SOURCE):

Type	PhD thesis
Title	Chemokine receptor signals: role in migration, invasion and cancer metastasis
Author(s)	I.S. Zeelenberg
Faculty	AMC-UvA
Year	2004

FULL BIBLIOGRAPHIC DETAILS:
http://hdl.handle.net/11245/1.222358

Copyright

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content licence (like Creative Commons).

UvA-DARE is a service provided by the library of the University of Amsterdam (http://dare.uva.nl)
(pagedate: 2014-11-16)

Contents

Chapter 1	Introduction	**9**
Chapter 2	Retention of CXCR4 in the endoplasmic reticulum blocks dissemination of a T cell hybridoma. *J. Clin. Invest.* 108:269-277 (2001)	**49**
Chapter 3	Stromal cell-derived factor-1-induced LFA-1 activation during in vivo migration of T cell hybridoma cells requires Gq/11, RhoA, and myosin, as well as Gi and Cdc42. *J. Immunol.* 166(7):4293-4301 (2001)	**61**
Chapter 4	Blocking G-protein βγ dimers reduces desensitization of CXCR4, causing enhanced and persistent migration, independent of PI3-kinase. *Submitted for publication*	**73**
Chapter 5	Fusion of docked vesicles, driven by synaptotagmin, is required for chemokine-induced T-cell migration. *To be submitted*	**85**
Chapter 6	The chemokine receptor CXCR4 is required for outgrowth of colon carcinoma micrometastases. *Cancer Res.* 63:3833-3839 (2003)	**97**
Chapter 7	A putative role for the chemokine receptor CXCR5 in tumor growth of colon carcinoma cells. *Preliminary report*	**107**
Summary		**117**
Samenvatting		**121**
Dankwoord		**127**
Curriculum vitae		**133**
List of publications		**135**