

File ID uvapub:89796  
Filename 336618.pdf  
Version final

---

SOURCE (OR PART OF THE FOLLOWING SOURCE):

Type article  
Title The CHESS spectral survey of star forming regions: Peering into the protostellar shock L1157-B1. I. Shock chemical complexity  
Author(s) C. Codella, B. Lefloch, C. Ceccarelli, J. Cernicharo, E. Caux, A. Lorenzani, S. Viti, P. Hily-Blant, B. Parise, S. Maret, B. Nisini, P. Caselli, et al.  
Faculty FNWI: Astronomical Institute Anton Pannekoek (IAP)  
Year 2010

FULL BIBLIOGRAPHIC DETAILS:

<http://hdl.handle.net/11245/1.336618>

---

*Copyright*

*It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content licence (like Creative Commons).*

---

LETTER TO THE EDITOR

# The CHESSE spectral survey of star forming regions: Peering into the protostellar shock L1157-B1

## I. Shock chemical complexity<sup>★,★★</sup>

C. Codella<sup>1</sup>, B. Lefloch<sup>2</sup>, C. Ceccarelli<sup>2</sup>, J. Cernicharo<sup>3</sup>, E. Caux<sup>4</sup>, A. Lorenzani<sup>1</sup>, S. Viti<sup>5,6</sup>, P. Hily-Blant<sup>2</sup>, B. Parise<sup>7</sup>, S. Maret<sup>2</sup>, B. Nisini<sup>8</sup>, P. Caselli<sup>9,1</sup>, S. Cabrit<sup>10</sup>, L. Pagani<sup>10</sup>, M. Benedettini<sup>6</sup>, A. Boogert<sup>11</sup>, F. Gueth<sup>12</sup>, G. Melnick<sup>13</sup>, D. Neufeld<sup>14</sup>, S. Pacheco<sup>2</sup>, M. Salez<sup>10</sup>, K. Schuster<sup>12</sup>, A. Bacmann<sup>2,15</sup>, A. Baudry<sup>15</sup>, T. Bell<sup>16</sup>, E. A. Bergin<sup>17</sup>, G. Blake<sup>16</sup>, S. Bottinelli<sup>4</sup>, A. Castets<sup>2</sup>, C. Comito<sup>7</sup>, A. Coutens<sup>4</sup>, N. Crimier<sup>2,3</sup>, C. Dominik<sup>18,19</sup>, K. Demyk<sup>4</sup>, P. Encrenaz<sup>10</sup>, E. Falgarone<sup>10</sup>, A. Fuente<sup>20</sup>, M. Gerin<sup>10</sup>, P. Goldsmith<sup>21</sup>, F. Helmich<sup>22</sup>, P. Hennebelle<sup>10</sup>, Th. Henning<sup>23</sup>, E. Herbst<sup>24</sup>, T. Jacq<sup>15</sup>, C. Kahane<sup>2</sup>, M. Kama<sup>18</sup>, A. Klotz<sup>2</sup>, W. Langer<sup>21</sup>, D. Lis<sup>16</sup>, S. Lord<sup>16</sup>, J. Pearson<sup>21</sup>, T. Phillips<sup>16</sup>, P. Saraceno<sup>6</sup>, P. Schilke<sup>7,25</sup>, X. Tielens<sup>26</sup>, F. van der Tak<sup>22</sup>, M. van der Wiel<sup>27,22</sup>, C. Vastel<sup>4</sup>, V. Wakelam<sup>15</sup>, A. Walters<sup>4</sup>, F. Wyrowski<sup>7</sup>, H. Yorke<sup>21</sup>, C. Borys<sup>16</sup>, Y. Delorme<sup>10</sup>, C. Kramer<sup>28</sup>, B. Larsson<sup>29</sup>, I. Mehdi<sup>21</sup>, V. Ossenkopf<sup>25</sup>, and J. Stutzki<sup>25</sup>

(Affiliations can be found after the references)

Received 30 March 2010 / Accepted 5 May 2010

### ABSTRACT

We present the first results of the unbiased survey of the L1157-B1 bow shock, obtained with HIFI in the framework of the key program Chemical *Herschel* Survey of Star forming regions (CHESSE). The L1157 outflow is driven by a low-mass Class 0 protostar and is considered the prototype of the so-called chemically active outflows. The bright blue-shifted bow shock B1 is the ideal laboratory for studying the link between the hot (~1000–2000 K) component traced by H<sub>2</sub> IR-emission and the cold (~10–20 K) swept-up material. The main aim is to trace the warm gas chemically enriched by the passage of a shock and to infer the excitation conditions in L1157-B1. A total of 27 lines are identified in the 555–636 GHz region, down to an average 3 $\sigma$  level of 30 mK. The emission is dominated by CO(5–4) and H<sub>2</sub>O(1<sub>10</sub>–1<sub>01</sub>) transitions, as discussed by Lefloch et al. in this volume. Here we report on the identification of lines from NH<sub>3</sub>, H<sub>2</sub>CO, CH<sub>3</sub>OH, CS, HCN, and HCO<sup>+</sup>. The comparison between the profiles produced by molecules released from dust mantles (NH<sub>3</sub>, H<sub>2</sub>CO, CH<sub>3</sub>OH) and that of H<sub>2</sub>O is consistent with a scenario in which water is also formed in the gas-phase in high-temperature regions where sputtering or grain-grain collisions are not efficient. The high excitation range of the observed tracers allows us to infer, for the first time for these species, the existence of a warm ( $\geq 200$  K) gas component coexisting in the B1 bow structure with the cold and hot gas detected from ground.

**Key words.** ISM: individual objects: L1157 – ISM: molecules – stars: formation

## 1. Introduction


A newborn protostar generates a fast and well collimated jet, possibly surrounded by a wider angle wind. In turn, the ejected material drives (bow-)shocks travelling through the surrounding high-density medium and traced by H<sub>2</sub> ro-vibrational lines at excitation temperatures of around 2000 K. As a consequence, slower and cold (10–20 K) molecular outflows are formed by swept-up material, usually traced by CO. Shocks heat the gas and trigger several processes such as endothermic chemical reactions and ice grain mantle sublimation or sputtering. Several molecular species undergo significant enhancements in their abundances (see e.g., van Dishoeck & Blake 1998), as observed by observations at millimeter wavelengths towards a number of outflows (Garay et al. 1998; Bachiller & Pérez Gutiérrez 1997, BP97 hereafter; Jørgensen et al. 2007). The link between the gas

components at ~10 K and the hot 2000 K shocked component is crucial to understanding how the protostellar wind transfers momentum and energy back to the ambient medium. In this context, the understanding of the chemical composition of a typical molecular bow-shock is essential because it represents a very powerful diagnostic tool for probing its physical conditions.

The L1157 outflow, located at a distance estimated to be between 250 pc (Looney et al. 2007) and 440 pc (Viotti 1969) may be regarded as the ideal laboratory for observing the effects of shocks on the gas chemistry, being the archetype of the so-called chemically rich outflows (Bachiller et al. 2001). The low-luminosity (4–11 L<sub>⊙</sub>) Class 0 protostar IRAS20386+6751 drives a precessing powerful molecular outflow associated with several bow shocks seen in CO (Gueth et al. 1996) and in IR H<sub>2</sub> images (Davis & Eislöffel 1995; Neufeld et al. 2009). In particular, the brightest blue-shifted bow-shock, called B1 (Fig. 1), has been extensively mapped with the PdB and VLA interferometers at mm- and cm-observations revealing a rich and clumpy structure, the clumps being located at the wall of the cavity with an arch-shape (Tafalla & Bachiller 1995; Gueth et al. 1998; Benedettini et al. 2007, hereafter BVC07; Codella et al. 2009).

\* *Herschel* is an ESA space observatory with science instruments provided by European-led principal Investigator consortia and with important participation from NASA.

\*\* Table 1 is only available in electronic form at <http://www.aanda.org>



**Fig. 1.** The B1 clump. PdBI emission of  $\text{CH}_3\text{OH}(2_{1-1_1})\text{A}^-$  (grey) on the CS(2–1) one (contours), from BVC07. The maps are centred on the coordinates used for the present HIFI observations  $\alpha_{J2000} = 20^{\text{h}}39^{\text{m}}10^{\text{s}}.2$ ,  $\delta_{J2000} = +68^{\circ}01'10''.5$ , i.e. at  $\Delta\alpha = +25''.6$  and  $\Delta\delta = -63''.5$  from the driving protostar. The labels indicate the main B1 clumps detected in different tracers. Circles are for the HPBWs of the HIFI data presented here (39'') and of Band 7 (11''), i.e., at the highest frequencies of the CHES surveys.

L1157-B1 is well traced by molecules thought to be released by dust mantles such as  $\text{H}_2\text{CO}$ ,  $\text{CH}_3\text{OH}$ , and  $\text{NH}_3$  as well as typical tracers of high-speed shocks such as SiO (e.g., Gusdorf et al. 2008). Temperatures  $\approx 60$ –200 K (from  $\text{NH}_3$ ,  $\text{CH}_3\text{CN}$ , and SiO) as well as around 1000 K (from  $\text{H}_2$ ) have been derived (Tafalla & Bachiller 1995; Codella et al. 2009; Nisini et al. 2007, in prep.). However, a detailed study of the excitation conditions of the B1 structure has yet to be completed because of the limited range of excitation covered by the observations performed so far at cm- and mm-wavelengths. Observations of sub-mm lines with high excitation ( $\geq 50$ –100 K above the ground state) are thus required.

As part of the *Herschel* key program CHES<sup>1</sup> (Chemical *HERschel* Surveys of Star forming regions), L1157-B1 is currently being investigated with an unbiased spectral survey using the HIFI instrument (de Graauw et al. 2010). In this Letter, we report the first results based on HIFI observations in the 555–636 GHz spectral window, confirming the chemical richness and revealing different molecular components at different excitation conditions coexisting in the B1 bow structure.

## 2. Observations

The observations were performed on 2009, August 1, during the Performance Verification phase of the HIFI heterodyne instrument (de Graauw et al. 2010) on board of the *Herschel* Space Observatory (Pilbratt et al. 2010). The band called 1b (555.4–636.2 GHz) was covered in double-sideband (DSB) with a total integration time of 140 min. The wide band spectrometer was used with a frequency resolution of 1 MHz. The typical HPBW is 39''. The data were processed with the ESA-supported package HIPE<sup>2</sup> (*Herschel* interactive processing environment) for baseline subtraction and sideband deconvolution and then analysed with the GILDAS<sup>3</sup> software. All the spectra (here in units of antenna  $T_a$ ) were smoothed to a velocity resolution of

1 km s<sup>-1</sup>, except those showing the weakest emission, which were smoothed to lower spectral resolutions (up to 4 km s<sup>-1</sup>). At a velocity resolution of 1 km s<sup>-1</sup>, the rms noise is 6–13 mK ( $T_a$  scale), depending on the line frequency. The main-beam efficiency ( $\eta_{\text{mb}}$ ) has not yet been reliably determined. When needed, we adopted an average  $\eta_{\text{mb}}$  of 0.72.

## 3. Different tracers at different velocities

A total of 27 emission lines were detected, with a wide range of upper level energies, from a few tens to a few hundreds of Kelvin. Table 1 lists the spectroscopic and observational parameters of all the transitions. For the first time, high excitation (up to  $\approx 200$  K) emission lines related to species whose abundance is largely enhanced in shocked regions were detected. The CO(5–4) and  $\text{H}_2\text{O}(1_{10-1_{01}})$  lines are analysed in Lefloch et al. (2010). Figure 2 presents representative examples of line profiles observed towards L1157-B1. All the spectra contain lines with blue-shifted wings peaking near 0 km s<sup>-1</sup>, which have a terminal velocity equal to  $\sim -8, -6$  km s<sup>-1</sup>. Previous PdBI observations showed that L1157-B1 is associated with very high velocities (HVs) of as low as  $\approx -20$  km s<sup>-1</sup> ( $v_{\text{LSR}} = +2.6$  km s<sup>-1</sup>, BP97). We cannot exclude the lack of detected emission in the HV regime in the present HIFI spectra being caused by their relatively low signal-to-noise (S/N) ratio. The PdBI images indicate that the brightness of the emission lines in the HV regime is indeed weaker than the emission at low velocities by a factor of 5–10. The spectra in Fig. 2 clearly show that this weak emission would lie below the noise. On the other hand, the HV gas is detected in the very bright lines of CO and  $\text{H}_2\text{O}$  (Lefloch et al. 2010). We note that the HV emission is mostly confined to within the eastern B1a clump (Fig. 1), within an emitting region of size  $\leq 10''$  (Gueth et al. 1998; BVC07), whereas low velocity lines originate in both the bow-structure and the walls of the outflow cavity (e.g., the B0e and B0d in Fig. 1), of typical size 15''–18''. Therefore, the forthcoming HIFI-CHES observations at higher frequencies and higher spatial resolution (see the dashed circle in Fig. 1) should allow us to study the HV wings in species other than CO and  $\text{H}_2\text{O}$ .


The uniqueness of HIFI lies in its high spectral profile resolution for many high excitation transitions of a large number of molecular species. The analysis of the present HIFI spectra reveals a secondary peak occurring between  $-3.0$  and  $-4.0$  km s<sup>-1</sup> (here defined medium velocity, MV) and well outlined by e.g., HCN(7–6). The MV peak is also visible in  $\text{NH}_3(1_{0-0_0})$  and in some lines of  $\text{CH}_3\text{OH}$  and  $\text{H}_2\text{CO}$  (see Fig. 3), but its occurrence does not show any clear trend with the choice of tracer of line excitation. No single-dish spectra had previously detected this spectral feature (BP97; Bachiller et al. 2001). An inspection of the spectra observed at PdBI shows that the MV secondary peak is observed in a couple of lines of the  $\text{CH}_3\text{OH}(2_{\text{K}}-1_{\text{K}})$  series (see Fig. 3 of BVC07) and only towards the western B1b clump (size  $\sim 5''$ ). This finding implies that there is a velocity component originating mainly in the western side of B1, while the HV gas is emitted from the eastern one (see above).

Figure 3 compares the profiles of the  $\text{NH}_3(1_{0-0_0})$  and  $\text{H}_2\text{CO}(8_{17-7_{16}})$  lines with the  $\text{H}_2\text{O}(1_{10-1_{01}})$  profile, where the S/N allows such an analysis (MV and LV ranges). By assuming that the emission in the MV range is optically thin (including the  $\text{H}_2\text{O}$  line) and originates in the same region, we obtained from the comparison of their profiles a straightforward estimate of the relative abundance ratios of the gas at different velocities. As a notable example, the  $\text{NH}_3/\text{H}_2\text{O}$  intensity ratio decreases by a factor  $\sim 5$  moving towards higher velocities (Fig. 3), implying


<sup>1</sup> <http://www-laog.obs.ujf-grenoble.fr/heberges/ches/>

<sup>2</sup> HIPE is a joint development by the *Herschel* Science Ground Segment Consortium, consisting of ESA, the NASA *Herschel* Science Center, and the HIFI, PACS and SPIRE consortia.

<sup>3</sup> <http://www.iram.fr/IRAMFR/GILDAS>



**Fig. 2.** Molecular line profiles observed towards L1157-B1: species and transitions are reported in the panels. The vertical solid line indicates the ambient LSR velocity ( $+2.6 \text{ km s}^{-1}$  from  $\text{C}^{18}\text{O}$  emission; BP97), while the dashed one is for the secondary peak at  $-4.0 \text{ km s}^{-1}$ .




**Fig. 3.** *Top and middle panels:* Comparison between the profiles of  $\text{NH}_3(1_0-0_0)$ , multiplied by a factor 7.4,  $\text{H}_2\text{CO}(8_{17}-7_{16})$ , multiplied by a factor 22.5,  $\text{HCN}(7-6)$ , multiplied by a factor 9.0, and  $\text{H}_2\text{O}(1_{10}-1_{01})$ , the latter from Lefloch et al. (2010). The vertical solid line indicates the ambient LSR velocity ( $+2.6 \text{ km s}^{-1}$ ). The velocity ranges arbitrarily defined as HV ( $-20, -6 \text{ km s}^{-1}$ ; traced by  $\text{H}_2\text{O}$ ), MV ( $-6, -1.5 \text{ km s}^{-1}$ ; outlined by the  $\text{HCN}$  and  $\text{H}_2\text{CO}$  secondary peak), and LV ( $-1.5, +2.6 \text{ km s}^{-1}$ ; the rest of the blue wing) are drawn (see text). *Bottom panel:* Intensity  $\text{NH}_3/\text{H}_2\text{O}$  line ratio as a function of velocity.

that a similar decrease in the abundance ratios occurs. This may reflect different pre-shock ice compositions in the gas emitting the MV emission. Alternatively, this behavior is consistent with  $\text{NH}_3$  being released by grain mantles, but water both being released by grain mantles and, in addition, copiously forming in the warm shocked gas from endothermic reactions, which convert all gaseous atomic oxygen into water (Kaufman & Neufeld 1997; Jiménez-Serra et al. 2008, and references therein). The water abundance may be enhanced with respect to ammonia in the fast and warm ( $\geq 220 \text{ K}$ ) gas, which might explain why the  $\text{H}_2\text{O}$  wings are larger than those of  $\text{NH}_3$ ,  $\text{CH}_3\text{OH}$ , and  $\text{H}_2\text{CO}$ , all species being directly evaporated from dust grain mantles.

#### 4. Physical properties along the B1 bow shock

We detected several lines from  $\text{CH}_3\text{OH}$  (17 lines with upper level energies up to 211 K). We can derive a first estimate of the emitting gas temperature by means of the standard analysis



**Fig. 4.** Rotation diagrams for the  $\text{CH}_3\text{OH}$  transitions measured with HIFI (triangles) and from ground (PdBI; squares). Black and blue points are for A- and E-form, respectively. The parameters  $N_u$ ,  $g_u$ , and  $E_u$  are, respectively, the column density, the degeneracy and the energy (with respect to the ground state of each symmetry) of the upper level. The derived values of the rotational temperature are reported: (i) 106 K, for the HIFI lines covering the  $E_u = 32\text{--}211 \text{ K}$  excitation range and (ii) 12 K (as Bachiller et al. 1995), for the PdBI lines, at lower excitation.

of the rotational diagram. We show the case of methanol (A- and E-forms) in Fig. 4. The derived rotational temperature ( $T_{\text{rot}}$ ) is 106 K (with an uncertainty of  $\sim 20 \text{ K}$ ), which represents a lower limit to the kinetic temperature ( $T_{\text{kin}}$ ). In the same figure, we report the methanol lines ( $2_K-1_K$ ) observed with PdBI and whose intensity is integrated in the HIFI 39'' beam. The  $T_{\text{rot}}$  derived from the ground-based data (based only on lines with  $E_u \leq 50 \text{ K}$ ; BVC07) is definitely lower,  $\sim 12 \text{ K}$ , in perfect agreement with that found with the 30-m spectra in the same excitation range by Bachiller et al. (1995). As discussed by Goldsmith & Langer (1999), this behavior may be caused by either two components at different temperatures or both non-LTE effects and line opacity. These two possibilities cannot be distinguished based only on the rotational diagram. However, given that a range of  $T_{\text{kin}}$  and  $n_{\text{H}_2}$  is naturally expected in a shock, if we were to assume that two gas components provide an explanation, they would not only have different temperatures but also a different column densities. Taking the filling factor  $f = 0.13$ , derived by the  $\text{CH}_3\text{OH}$  maps obtained at the PdBI, the low temperature component column density is  $8 \times 10^{14} \text{ cm}^{-2}$  (in agreement with Bachiller et al. 1995), whereas the high temperature component has a column density of around  $10^{14} \text{ cm}^{-2}$ . We note that the rotation diagrams

obtained for the MV and LV CH<sub>3</sub>OH emission separately do not allow us to infer any clear difference.

It is possible to more tightly constrain the emitting gas temperature and  $n_{\text{H}_2}$  density for the species where the collisional rate coefficients are known, by performing a non-LTE analysis. To this end, we used the non-LTE excitation code RADEX with an escape probability formalism for the radiative transfer (Van der Tak et al. 2007) coupled with the LAMDA database (Schöier et al. 2005). Methanol is the species detected in the largest number of lines. The full non-LTE study will be reported in a forthcoming paper. Here we analysed only the E-form, for which the collisional rate coefficients are available (Pottage et al. 2004). The major result of this analysis is that for a range of densities of  $10^3$ – $10^7$  cm<sup>-3</sup>, the gas temperature exceeds 200 K. A similar result is obtained by considering H<sub>2</sub>CO emission.

Finally, by combining the HIFI CS(12–11) line with CS(2–1) and (3–2) lines observed with ground-based telescopes, we also derive a kinetic temperature that is definitely above 300 K for the outflowing gas. In this case, caution should be taken since we are able to trace different gas components, as suggested by CH<sub>3</sub>OH, the gas at higher excitation being traced by CS(12–11). If we analyse only the (2–1)/(3–2) intensity ratio, the non-LTE approach does not allow us to constrain the temperature in this way, but we are able to infer  $n_{\text{H}_2}$  of around  $4 \times 10^4$  cm<sup>-3</sup>. Interestingly, when we check for a possible dependence of  $n_{\text{H}_2}$  on velocity, the LV range is found to be indicative of a denser medium ( $\sim 10^5$  cm<sup>-3</sup>) by an order of magnitude with respect to the MV gas.

## 5. Conclusions

We have presented the HIFI unbiased spectral survey in the 555–636 GHz band towards the bright bow-shock B1 of the L1157 protostellar outflow. For the first time, we have detected high-excitation (up to  $\approx 200$  K) emission lines of species whose abundance is largely enhanced in shocked regions (e.g., H<sub>2</sub>O, NH<sub>3</sub>, H<sub>2</sub>CO, CH<sub>3</sub>OH). This has allowed us to trace with these species the existence of a high excitation component with  $T_{\text{kin}} \geq 200$ –300 K. Temperature components from  $\sim 300$  K to  $\sim 1400$  K have been inferred from the analysis of the H<sub>2</sub> pure rotational lines (Nisini et al., in prep.). Therefore the present observations provide a link between the gas at  $T_{\text{kin}} 60$ –200 K previously observed from the ground and the warmer gas probed by the H<sub>2</sub> lines. We plan to perform additional HIFI observations in the THz region towards L1157-B1 to observe more species and transitions, thus to be able to derive reliable abundances and study of the different gas components associated with the bow structure.

*Acknowledgements.* HIFI has been designed and built by a consortium of institutes and university departments from across Europe, Canada and the United States under the leadership of SRON Netherlands Institute for Space Research, Groningen, The Netherlands and with major contributions from Germany, France and the US. Consortium members are: Canada: CSA, U. Waterloo; France: CESR, LAB, LERMA, IRAM; Germany: KOSMA, MPIfR, MPS; Ireland: NUI Maynooth; Italy: ASI, IFSI-INAF, Osservatorio Astrofisico di Arcetri-INAF; Netherlands: SRON, TUD; Poland: CAMK, CBK; Spain: Observatorio Astronómico Nacional (IGN), Centro de Astrobiología (CSIC-INTA). Sweden: Chalmers University of Technology – MC2, RSS & GARD; Onsala Space Observatory; Swedish National Space Board, Stockholm University – Stockholm Observatory; Switzerland: ETH Zurich, FHNW; USA: Caltech, JPL, NHSC. We thank many funding agencies for financial support.

## References

Bachiller, R., & Pérez Gutiérrez, M. 1999, *ApJ*, 487, L93 (BP97)  
 Bachiller, R., Liechti, S., Walmsley, C. M., & Colomer, F. 1995, *A&A*, 295, L51  
 Bachiller, R., Pérez Gutiérrez, M., Kumar, M. S. N., & Tafalla, M. 2001, *A&A*, 372, 899

Benedettini, M., Viti, S., Codella, C., et al. 2007, *MNRAS*, 381, 1127 (BVC07)  
 Codella, C., Benedettini, M., Beltrán, M. T., et al. 2009, *A&A*, 507, L25  
 Davis, C. J., & Eisloffel, J. 1995, *A&A*, 300, 851  
 de Graauw, Th., Helmich, F. P., Philipps, T. G., et al. 2010, *A&A*, 518, L6  
 Garay, G., Köhnenkamp, I., Bourke, T. L., Rodríguez, L. F., & Lehtinen, K. K. 1998, *ApJ*, 509, 768  
 Goldsmith, P. F., & Langer, W. D. 1999, *ApJ*, 517, 209  
 Gueth, F., Guilloteau, S., & Bachiller, R. 1996, *A&A*, 307, 891  
 Gueth, F., Guilloteau, S., & Bachiller, R. 1998, *A&A*, 333, 287  
 Gusdorf, A., Pineau Des Forêts, G., Cabrit, S., & Flower, D. R. 2008, *A&A*, 490, 695  
 Jiménez-Serra, I., Caselli, P., Martín-Pintado, J., & Hartquist, T. W. 2008, *A&A*, 482, 549  
 Jørgensen, J. K., Bourke, T. L., & Myers, P. C. 2007, *ApJ*, 659, 479  
 Lefloch, B., Cabrit, S., Codella, C., et al. 2010, *A&A*, 518, L113  
 Looney, L. W., Tobin, J. J., & Kwon, W. 2007, *ApJ*, 670, L131  
 Müller, H. S. P., Schöier, F. L., Stutzki, J., & Winnewisser, G. 2005, *J. Mol. Struct.*, 742, 215  
 Neufeld, D. A., Nisini, B., Giannini, T., et al. 2009, *ApJ*, 706, N09  
 Nisini, B., Codella, C., Giannini, T., et al. 2007, *A&A*, 462, 163  
 Pickett, H. L., Poynter, R. L., Cohen, E. A., et al. 1998, *J. Quant. Spectrosc. Rad. Transfer*, 60, 883  
 Pilbratt, G. L., Riedinger, J. L., Passvogel, T., et al. 2010, *A&A*, 518, L1  
 Pottage, J. T., Flower, D. R., & Davis, S. L. 2004, *MNRAS*, 352, 39  
 Schöier, F. L., van der Tak, F. F. S., van Dishoeck, E. F., & Black, J. H. 2005, *A&A*, 432, 369  
 Tafalla, M., & Bachiller, R. 1995, *ApJ* 443, L37  
 Viotti, N. R. 1969, *Mem. Soc. Astron. Ital.*, 40, 75  
 van Dishoeck, E. F., & Blake, G. A. 1998, *ARA&A*, 36, 317  
 Van der Tak, F. F. S., Black, J. H., Schöier, F. L., Jansen, D. J., & van Dishoeck, E. F. 2007, *A&A*, 468, 627

- <sup>1</sup> INAF, Osservatorio Astrofisico di Arcetri, Firenze, Italy  
e-mail: codella@arcetri.astro.it
- <sup>2</sup> Laboratoire d'Astrophysique de Grenoble, UMR 5571-CNRS, Université Joseph Fourier, Grenoble, France
- <sup>3</sup> Centro de Astrobiología, CSIC-INTA, Madrid, Spain
- <sup>4</sup> CESR, Université Toulouse 3 and CNRS, Toulouse, France
- <sup>5</sup> Department of Physics and Astronomy, University College London, London, UK
- <sup>6</sup> INAF, Istituto di Fisica dello Spazio Interplanetario, Roma, Italy
- <sup>7</sup> Max-Planck-Institut für Radioastronomie, Bonn, Germany
- <sup>8</sup> INAF, Osservatorio Astronomico di Roma, Monte Porzio Catone, Italy
- <sup>9</sup> School of Physics and Astronomy, University of Leeds, Leeds, UK
- <sup>10</sup> Observatoire de Paris-Meudon, LERMA UMR CNRS 8112, Meudon, France
- <sup>11</sup> Infrared Processing and Analysis Center, Caltech, Pasadena, USA
- <sup>12</sup> Institut de RadioAstronomie Millimétrique, Grenoble, France
- <sup>13</sup> Center for Astrophysics, Cambridge MA, USA
- <sup>14</sup> Johns Hopkins University, Baltimore MD, USA
- <sup>15</sup> CNRS/INSU, Laboratoire d'Astrophysique de Bordeaux, Floirac, France
- <sup>16</sup> California Institute of Technology, Pasadena CA, USA
- <sup>17</sup> University of Michigan, Ann Arbor, MI 48109, USA
- <sup>18</sup> Astronomical Institute Anton Pannekoek, University of Amsterdam, Amsterdam, The Netherlands
- <sup>19</sup> Department of Astrophysics/IMAPP, Radboud University Nijmegen, Nijmegen, The Netherlands
- <sup>20</sup> IGN Observatorio Astronómico Nacional, Alcalá de Henares, Spain
- <sup>21</sup> Jet Propulsion Laboratory, Caltech, Pasadena, CA 91109, USA
- <sup>22</sup> SRON, Institute for Space Research, Groningen, The Netherlands
- <sup>23</sup> Max-Planck-Institut für Astronomie, Heidelberg, Germany
- <sup>24</sup> Ohio State University, Columbus OH, USA
- <sup>25</sup> Physikalisches Institut, Universität zu Köln, Köln, Germany
- <sup>26</sup> Leiden Observatory, Leiden University, Leiden, The Netherlands
- <sup>27</sup> Kapteyn Astronomical Institute, Groningen, The Netherlands
- <sup>28</sup> Institut de RadioAstronomie Millimétrique, Granada, Spain
- <sup>29</sup> Department of Astronomy, Stockholm University, Stockholm, Sweden

**Table 1.** List of molecular species and transitions observed with HIFI (Band 1b).

Transition	$\nu_0^a$ (MHz)	$E_u^a$ (K)	$T_{\text{peak}}$ (mK)	rms (mK)	$V_{\text{peak}}$ (km s <sup>-1</sup> )	$V_{\text{min}}$ (km s <sup>-1</sup> )	$V_{\text{max}}$ (km s <sup>-1</sup> )	$F_{\text{int}}$ (K km s <sup>-1</sup> )
o-H <sub>2</sub> O(1 <sub>10</sub> -1 <sub>01</sub> )	556 936.002	27	910(17)	17	-0.37(1.00)	-25.4	+7.6	11.68(0.10)
CH <sub>3</sub> OH E (11 <sub>2,9</sub> - 10 <sub>1,9</sub> )	558 344.500	168	47(10)	10	+0.60(1.00)	-2.9	+2.7	0.16(0.02)
o-H <sub>2</sub> CO(8 <sub>18</sub> -7 <sub>17</sub> )	561 899.318	118	92(4) <sup>b</sup>	8	+0.38(0.14) <sup>b</sup>	-4.5	+3.6	0.49(0.03) <sup>b</sup>
CH <sub>3</sub> OH E (3 <sub>2,2</sub> - 2 <sub>1,2</sub> )	568 566.054	32	42(8)	8	+0.60(1.00)	-2.6	+2.7	0.31(0.02)
o-NH <sub>3</sub> (1 <sub>0</sub> -0 <sub>0</sub> )	572 498.068	28	122(7)	7	+1.03(0.80)	-6.9	+5.0	0.89(0.03)
CO(5-4)	576 267.931	83	883(10)	10	+1.60(1.00)	-36.5	+6.0	49.30(0.07)
p-H <sub>2</sub> CO(8 <sub>08</sub> -7 <sub>07</sub> )	576 708.315	125	41(7)	7	+1.80(1.00)	-5.4	+2.7	0.22(0.02)
CH <sub>3</sub> OH A <sup>-</sup> (2 <sub>2,1</sub> - 1 <sub>1,0</sub> )	579 084.700	45	53(8)	8	+0.60(1.00)	-6.0	+3.7	0.29(0.03)
CH <sub>3</sub> OH E (12 <sub>1,12</sub> - 11 <sub>1,11</sub> )	579 151.003	178	44(8)	8	-0.30(1.00)	-4.0	+4.0	0.25(0.02)
CH <sub>3</sub> OH A <sup>+</sup> (12 <sub>0,12</sub> - 11 <sub>0,11</sub> )	579 459.639	181	48(7)	7	+0.90(1.00)	-3.9	+4.9	0.27(0.02)
CH <sub>3</sub> OH A <sup>+</sup> (2 <sub>2,0</sub> - 1 <sub>1,1</sub> )	579 921.342	45	42(7)	7	-0.60(1.00)	-3.3	+2.6	0.21(0.02)
CH <sub>3</sub> OH E (12 <sub>2,10</sub> - 11 <sub>2,9</sub> )	580 902.721	195	11(4)	4	-3.00(3.00)	-3.9	+2.5	0.09(0.02)
CH <sub>3</sub> OH A <sup>+</sup> (6 <sub>1,6</sub> - 5 <sub>0,5</sub> )	584 449.896	63	88(9)	9	-0.30(1.00)	-6.0	+2.9	0.58(0.03)
CS(12-11)	587 616.240	183	23(2) <sup>b</sup>	5	-0.61(0.57) <sup>b</sup>	-7.3	+6.5	0.19(0.03) <sup>b</sup>
CH <sub>3</sub> OH A <sup>+</sup> (7 <sub>3,5</sub> - 6 <sub>2,4</sub> )	590 277.688	115	42(9)	9	+0.60(1.00)	-1.0	+3.0	0.16(0.02)
CH <sub>3</sub> OH A <sup>-</sup> (7 <sub>3,4</sub> - 6 <sub>2,5</sub> )	590 440.291	115	40(9)	9	-0.60(1.00)	-2.6	+0.8	0.10(0.02)
CH <sub>3</sub> OH E (9 <sub>0,9</sub> - 8 <sub>1,8</sub> )	590 790.957	110	40(10)	10	+0.60(1.00)	-4.7	+4.3	0.28(0.03)
o-H <sub>2</sub> CO(8 <sub>17</sub> -7 <sub>16</sub> )	600 374.604	126	29(7) <sup>b</sup>	9	-0.14(0.57) <sup>b</sup>	-3.0	+1.8	0.19(0.04) <sup>b</sup>
CH <sub>3</sub> OH E (4 <sub>2,3</sub> - 3 <sub>1,3</sub> )	616 979.984	41	47(6)	6	+0.90(1.00)	-4.5	+2.6	0.26(0.02)
HCN(7-6)	620 304.095	119	94(7)	7	-0.60(1.00)	-7.6	+3.2	0.68(0.03)
HCO <sup>+</sup> (7-6)	624 208.180	119	30(3) <sup>b</sup>	8	+0.53(0.47) <sup>b</sup>	-3.6	+4.5	0.11(0.03) <sup>b</sup>
CH <sub>3</sub> OH A <sup>-</sup> (3 <sub>2,2</sub> - 2 <sub>1,1</sub> )	626 626.302	52	18(5)	5	-1.20(4.00)	-2.5	+6.3	0.18(0.07)
CH <sub>3</sub> OH E (13 <sub>1,13</sub> - 12 <sub>1,12</sub> )	627 170.503	209	33(7)	7	-0.30(1.00)	-3.2	+2.5	0.15(0.02)
CH <sub>3</sub> OH A <sup>+</sup> (13 <sub>0,13</sub> - 12 <sub>0,12</sub> )	627 558.440	211	41(9)	9	+0.60(1.00)	-3.6	+2.5	0.19(0.02)
CH <sub>3</sub> OH A <sup>+</sup> (3 <sub>1,2</sub> - 2 <sub>1,2</sub> )	629 140.493	52	52(9)	9	+0.60(1.00)	-3.5	+3.7	0.24(0.03)
CH <sub>3</sub> OH A <sup>+</sup> (7 <sub>1,7</sub> - 6 <sub>0,6</sub> )	629 921.337	79	70(13)	13	+1.50(1.00)	-3.9	+3.7	0.37(0.04)
o-H <sub>2</sub> CO(9 <sub>19</sub> -8 <sub>18</sub> )	631 702.813	149	66(4) <sup>b</sup>	9	+0.45(0.17) <sup>b</sup>	-1.5	+2.6	0.22(0.02) <sup>b</sup>

**Notes.** CO and H<sub>2</sub>O emission is discussed in Lefloch et al. (2010). Peak velocity and intensity (in  $T_a$  scale), integrated intensity ( $F_{\text{int}}$ ), as well as the terminal velocities of the line emission ( $V_{\text{min}}$  and  $V_{\text{max}}$ ) are reported. <sup>(a)</sup> Frequencies and spectroscopic parameters have been extracted from the Jet Propulsion Laboratory molecular database (Pickett et al. 1998) for all the transition except those of CH<sub>3</sub>OH, which have been extracted from the Cologne Database for Molecular Spectroscopy (Müller et al. 2005). Upper level energies refer to the ground state of each symmetry. <sup>(b)</sup> Gaussian fit.