

Is the Extremely Low Heterozygosity Level in Yponomeuta rorellus Caused by Bottlenecks?

Steph B. J. Menken

Evolution, Vol. 41, No. 3. (May, 1987), pp. 630-637.

Stable URL:

http://links.jstor.org/sici?sici=0014-3820%28198705%2941%3A3%3C630%3AITELHL%3E2.0.CO%3B2-Z

Evolution is currently published by Society for the Study of Evolution.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at http://www.jstor.org/about/terms.html. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at http://www.jstor.org/journals/ssevol.html.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

IS THE EXTREMELY LOW HETEROZYGOSITY LEVEL IN *YPONOMEUTA***RORELLUS CAUSED BY BOTTLENECKS?

STEPH B. J. MENKEN

Department of Botany, University of Amsterdam, Kruislaan 318, 1098 SM Amsterdam, The Netherlands

Abstract.—Among the common and widespread species of the small ermine moths (Yponomeuta) the almost complete absence of genetic variability from a sample of 75 protein loci is a striking feature of Y. rorellus. Congeners exhibit normal to high H levels. At least for soluble proteins, this variation estimate is a real one, not resulting from an inability of the zymogram technique to detect variation nor from a biased sample of loci. The most likely explanation for the dearth of variation observed is a bottleneck at the species' origin. Moreover, ongoing population-size fluctations in the historical past of Y. rorellus may have prevented the recovery of equilibrium H levels as a consequence of genetic drift in small populations.

Received October 2, 1985. Accepted November 24, 1986

Genetic variation within populations is a prerequisite for evolution, as it provides the raw material for selection of better-adapted genotypes in response to challenges by the environment. Mutation is the ultimate source of all this variation.

Rapidly accumulating information from the last two decades shows that many species have high levels of genetic variation at the enzyme level; there is, however, much heterogeneity in variability levels among taxonomic groups as well as among species within taxonomic groups (Nevo et al., 1984; Graur, 1985).

Much speculation has arisen concerning the causes of these differences in overall genetic diversity. The following potential causes have been proposed: differences in population size, differences in environmental heterogeneity, genetic or breeding systems limiting variability, directional selection for homozygosity or for general-purpose genotypes in some species, or any combination of the aforementioned factors (Futuyma, 1979; Kimura, 1982; Nevo et al., 1984). A confusing factor in comparisons of heterozygosity estimates among species can be different samples of loci, as single locus heterozygosities differ much among loci (Powell, 1975; Menken, 1980c; Simon and Archie, 1985).

Species of the small ermine moth genus *Yponomeuta* (Lepidoptera, Yponomeutidae) have similar life histories, morphologies, and geographic distributions, but somewhat differing feeding habits (Herrebout et al., 1976). In a group of six closely

related species, heterozygosity levels per population based upon a homologous set of 51 genetic loci range from as low as 0% (Y. rorellus) to over 15% (Y. evonymellus, Y. padellus, and Y. cagnagellus; Table 4). In the present study, I report the near absence of any genetic variation in natural populations of Y. rorellus based upon a sample of 75 loci coding for soluble proteins. The reliability of the variation estimates and the possible causes of the low heterozygosity values will be discussed. Bottlenecks seem to be the major factor that account for the observed monomorphism.

Materials and Methods Life Cycle

Yponomeuta rorellus, a diploid, bisexual, univoltine species, is common and widespread with a palearctic distribution. Eggs are laid in July in masses on branches of willow (Salix spp.). Although the caterpillars hatch some three weeks after oviposition, they remain under the protection of the egg mass until the following spring. They then spin a communal web over the developing foliage and feed within or in the neighborhood of the web, extending it as the leaves are consumed. Entire willow trees may be defoliated in cases of severe infestation, a characteristic Y. rorellus shares with the other species of the so-called "padellus complex." In case of depletion of willows, few if any other hosts will be attacked (Herrebout et al., 1976). Fifth-instar full-grown caterpillars usually pupate within their web.



Fig. 1. Collection sites in the Netherlands for the *Yponomeuta rorellus* populations studied: 1) Haastrecht; 2) Hazerswoude; 3) Oosthuizen; 4) Schuilenburg (Kesteren); 5) Ter Aar; 6) Wageningen; 7) Zuidoost Beemster; 8) Zoetermeer.

Eclosion takes place after approximately three weeks.

Samples

Over a period of seven years (1974–1980) populations of *Y. rorellus* were sampled by collecting third- or fourth-instar larvae from the field (Table 1). They were reared in cages in the laboratory until eclosion. Mortality in the laboratory was less than 5%. Adults and, in some cases, fifth-instar larvae were deep frozen at -30°C until used for electrophoresis. Eight localities were sampled (Fig. 1). Food plants were *Salix alba* (most cases), *S. cinerea*, *S. viminalis*, *S. dasyclados*, *S. fragilis*, and *S. caprea*. Other host plants include *S. tenuijulis*, *S. babylonica*, *S. incana*, *S. oxycarpa*, and *S. australior* (Z. Gershenson, pers. comm.).

Electrophoresis

Specimen preparation, electrophoresis, isozyme specification, and enzyme assays follow Menken (1982a). In addition to the 51 loci studied in Menken (1982a) (Oosthuizen population, sampled in 1978), data are presented here on 22 more loci (Table

2). This set of loci was not included in the previous study because of inconsistently scorable variation and/or absence in one or more of the other small ermine moths investigated.

The variation reported is inferred to be Mendelian from comparison with the results from crosses in *Y. cagnagellus* (Menken, 1980a) and the general fit of observed phenotypic proportions to those expected under Hardy-Weinberg equilibrium.

The one heterozygote that occurs at each of the Fudh and $Est\beta$ -1 loci is a single male. This observation is in agreement with the sex-linked character of these two loci (Menken, 1980a; Lepidoptera have female heterogamety). A-Acph activity is limited to males (Menken, 1982a).

RESULTS

Natural populations of Y. rorellus harbor extremely low levels of genetic variation as revealed by electrophoresis. Individual values of mean heterozygosity per population (H) range from 0 to 0.0153 (mean \pm SE = 0.0045 \pm 0.0042; Table 1). Most notable is the Schuilenburg population, occurring on the common food plant S. alba, in which not a single individual in a sample of 25 moths was heterozygous for any of 45 proteins screened.

Glyceraldehyde-3-phosphate dehydrogenase contributes most to H; out of 105 mutant alleles, 58 or 55.2% are allele $G3pdh^{96}$ (Table 1). This allele is unique to Y. rorellus (Table 3). Single-locus heterozygosity ranges from 0.0000 to 0.4442 with a mean of 0.1103 \pm 0.1416. In contrast, G3pdh is extremely monomorphic in nine other Yponomeuta species (Menken, 1980c); just five heterozygotes were found in 7,000 individuals).

At 15 other loci, one or a few heterozygotes were found (Table 1). Every locus has one single mutant allele except for Pgm (alleles 97 and 103) and Aph-2 (alleles 94 and 103). Many of these alleles are unique to Y. rorellus (Table 3). Variant alleles at the $Est\beta-3$ (see below), Lap-3, and Lap-4 loci could not be classified as unique or otherwise, since they have not been studied in other Yponomeuta species.

Scarce information is available on two more loci not shown in Tables 1 and 2, an

								Heter	Heterzygotes		
Location	Food plant	Year	$N_{ m I}$	$N_{\rm L}$	Н	G3pdh			Other		
Adults:											
Ter Aar	Salix alba	1974	20	12	0.0049	1	3 Pgm				
Ter Aar	Salix alba	1975	20	74	0.0041	1	1 Pgm	1 Fudh	3 Aph-2		
Ter Aar	Salix alba	1976	20	40	0.0048	4		1 Lap-3	1 Aph-2		
Ter Aar	Salix alba	1977	20	20	0.0043	7	2 Pgm	2 A-6Pgdh	1 Fudh	2 Aph-2	2 Got-2
Ter Aar	Salix alba	1978	20	46	0.0033	ĸ	3 A-6Pgdh	1 $Est\beta$ -I	1 Lap-3		
Oosthuizen	Salix alba	1977	20	49	0.0036	0	2 A-Gluo	1 Mdh-2	1 Pt-1	3 Aph-2	2 Lap-1
Oosthuizen	Salix alba	1978	39	49a	0.0032	ĸ	1 Aph-2	1 Pt-1	1 A-Gluo	1 Lap-4	•
Oosthuizen	Salix alba	1980	18	31	0.0034	7					
Haastrecht	Salix alba	1978	25	4	0.0009	0	1 A-Acph				
Haastrecht	Salix alba	1979	4	32	0.0050	7					
Haastrecht	Salix cinerea	1979	16	30	0.0114	7					
Haastrecht	Salix viminalis	1979	16	30	0.0000	0					
Haastrecht	Salix dasyclados	1979	25	34	0.0106	10	1 A-Ao				
Wageningen	Salix alba	1977	20	47	0.0008	0	1 Pgm	1 Lap-4			
Schuilenburg	Salix alba	1978	25	45	0.0000	0					
ZO Beemster	Salix alba	1980	70	31	0.0074	4	1 Lap-4				
Hazerswoude	Salix fragilis	1979	18	59	9600.0	9					
Zoetermeer	Salix caprea	1979	15	59	0.0153	∞					
Larvae:											
Oosthuizen	Salix alba	1975	11	18	0.0000	0					
Oosthuizen	Salix alba	1978	25	21^{b}	0.0000	0					
Ter Aar	Salix alba	1977	25	22	0.0018	0	1 L- $Gluo$				

^a Based on 36 loci from Menken (1982a) plus 13 loci not studied therein. ^b Based on 15 loci from Menken (1982a) plus 6 loci not studied therein. ^c One homozygote for allele $G3pdh^{96}$ was also observed.

Table 2. Enzyme assays and stage specifications of Y. rorellus. See Menken (1982a) for composition of buffers symbolized. Symbols for quaternary structure: M = monomer; D = dimer; - = unknown. Stage specifications: Id = identical isozymes in both stages; Ss = stage specific (referred to as L and A forms); Ls = larva specific; As = adult specific; - = studied in adults only.

Enzyme (EC number)	Abbreviation for locus	Buffera	Quaternary structure	Stage speci- fication
Oxidoreductases				
Succinate dehydrogenase (1.3.99.1)	Sdh	I	_	Id
Glutamate dehydrogenase (1.4.1.2)	Gdh	I		_
Monoamine oxidase (1.4.3.4)	Mao	I	_	_
Tyrosinase (1.14.18.1)	L- Tyr	II	_	Ssa
Transferases				
Glutamic oxaloacetic transaminase (2.6.1.1)	Got-1	III		Id ^b
Hexokinase (2.7.1.1)	Hk-2	I		Id
Adenylate kinase (2.7.4.3)	Ak	II	_	Id
Hydrolases				
Esterase (3.1.1.2)	Esta-3	III	_	As
Esterase (3.1.1.2)	$Est\beta$ -2	III		As
Esterase (3.1.1.2)	Estβ-4	III	_	Ls
Alkaline phosphatase (3.1.3.1)	Aph-1	II	_	Ls
Alkaline phosphatase (3.1.3.1)	Aph-3	II	_	As
Amylase (3.2.1.1)	Amy-2	Veronal	_	Ls
β -glucosidase (3.2.1.21)	β -gluc	II	_	Ls
Leucine aminopeptidase (3.4.11.1)	Lap-2	I	_	Ls
Leucine aminopeptidase (3.4.11.1)	Lap-3	I	D	As
Leucine aminopeptidase (3.4.11.1)	Lap-4	I	M	As
Lyases				
Fumarase (4.2.1.2)	Fum	I		Id
Aconitase (4.2.1.3)	Acon	I	_	Id
Isomerases				
Triosephosphate isomerase (5.3.1.1)	Tpi	I	_	_
Other systems				
General protein	Pt-6	II	_	Id
General protein	Pt-7	II	_	Id

^a Adult form studied in Menken (1982a).

esterase ($Est\beta$ -3, EC 3.1.1.2) and mannose phosphate isomerase (Mpi, EC 5.3.1.8). Both code for monomeric enzymes. Detection of $Est\beta$ -3 is restricted to fresh adult material, and Mpi has been studied only once (Ter Aar population in 1981). The two enzymes seem to exhibit variation levels comparable with G3pdh.

The Ter Aar and Oosthuizen populations show quite constant H values over years and clearly have the highest amount of variation at loci other than G3pdh (Table 1). Variation at the A-6Pgdh locus (not investigated in 1974 and 1975) is restricted to Ter Aar. In general the other populations are somewhat more variable ($\bar{H}=0.0061\pm0.0055$ versus $\bar{H}=0.0030\pm0.0017$ for Ter Aar and Oosthuizen; two-tailed t-test, ns), due to the presence of a higher number of G3pdh

heterozygotes. In Ter Aar and Oosthuizen, a slightly smaller number of loci have been investigated on the average ($\bar{N} = 32.9$), compared with the other populations together $(\bar{N} = 34.8)$; hence no influence on H is to be expected (Nei and Roychoudhury, 1974; Graur, 1985). Those enzymes that appear to be polymorphic (1% criterion) in Ter Aar and Oosthuizen have been included in most other population surveys, where they appear to be monomorphic. A much more pronounced difference in H can be found between populations infesting S. alba, on the one hand, and all other Salix spp. lumped, on the other (Table 1; \bar{H} = 0.0030 ± 0.0022 and 0.0094 ± 0.0057 respectively; P < 0.01, two-tailed t-test). Again there is not much difference in the number of loci studied ($\bar{N} = 34.9$ and 30.4 respec-

b Run cathodally

tively). The lower value for H on S. alba can be fully ascribed to a lower incidence of variation at the G3pdh locus, which is not compensated by more variation at loci other than G3pdh (see above). Population sizes are comparable on the various food plants (W. Ravensbergen, pers. comm.).

Esterases are known to be among the most polymorphic enzymes; however, $Est\beta-3$ seems to be the only enzyme with some variation among seven isozyme loci.

DISCUSSION

Without genetic variability there can be no response to selection or genetic drift, and evolution cannot occur. The best single estimate of genetic variation in a population is the expected frequency of heterozygous loci per individual (H), assuming Hardy-Weinberg equilibrium (Nei and Roychoudhury, 1974). As these authors have pointed out, it is better to examine a large number of loci rather than a large number of individuals per locus for the reliable estimation of average heterozygosities (Archie, 1985). However, a negative correlation may exist between the number of enzymes surveyed and the magnitude of H, as studies often start with known polymorphic loci (Graur, 1985). Among a sample of 73 genetic loci, only G3pdh occasionally reaches appreciable variation levels in natural populations of Y. rorellus, resulting in H values up to 0.0153, based on various numbers of loci (Table 1).

Are these estimates an artifact caused by inadequate resolving power of the zymogram technique or biased sampling? It is now evident that electrophoresis is an imperfect detector of enzyme variants (heterogeneity within electromorphs: King and Ohta, 1975; Ayala, 1982). In order to elucidate part of this cryptic variation, some enzyme systems in Y. rorellus were subjected to sequential electrophoresis using two concentrations of starch (10 and 12%) and a total of 11 buffer systems with pH values ranging from 5.8 to 9.0 (unpubl.; see also Menken, 1982a). No additional variation was found at the G3pdh, Pgi, Pgm, Lap-1, $Est\beta-1$, $Est\beta-2$, and A-Gluo loci, loci that are known (with the exception of G3pdh) to be highly polymorphic in related Yponomeuta species (Menken, 1980c, 1981, 1982a). Thus the low level of variability in

Table 3. Alleles unique to Y. rorellus are designated by +; those designated by - also occur in congeners. All loci have 100 as the most-common allele (which is also the most-common allele in Y. cagnagellus [Menken, 1982a]), with the exceptions of five loci (*) that lack such alleles and which are diagnostic, and A-6Pgdh, Pt-1, and Mpi (with most-common alleles = 98, 98, and 95, respectively). The diagnostic loci are fixed save for A-Ao (most-common allele = 98) and L-Gluo (most-common allele = 104). For abbreviations see Table 2 and Menken (1982a).

Locus	Allele	Unique
Fudh	97	+
G3pdh	96	+
Idh-1*	104	+
Mdh-2	106	_
L-6Pgdh*	94	+
A-6Pgdh	97	+
A-Ao*	95	+
A-Ao*	98	+
A-Gluo	103	_
L-Gluo*	104	+
L-Gluo*	106	+
Got-2	98	+
Pgm	97	_
Pgm	103	_
Estα-1*	96	+
Estβ-1	104	+
A-Acph	98	_
Aph-2	94	+
Aph-2	103	_
Lap-1	107	+
Mpi	89	+
Pt-1	97	

Y. rorellus seems to be a real phenomenon, not caused by inadequate technical separation of proteins. Moreover, similar electrophoretic surveys in this laboratory have found normal H levels in several other insect species (Menken, 1982b; Menken and Ulenberg, 1983, 1986; Wilkinson et al., 1983).

Bearing the above in mind, one may conclude that *Y. rorellus* is extremely monomorphic at loci coding for soluble proteins, while all species closely related to *Y. rorellus* show mean heterozygosity levels ranging from 6 to 15% (Table 4). How can this low level of polymorphism be explained?

Restricted distribution and apomixis can be excluded as possible factors reducing variation, because this is a common, widespread species which should harbor a lot of variation according to the theoretical considerations of Kimura and Crow (1964) (see Varvio-Aho [1981] for a practical example). *Y. rorellus* has a normal sex ratio (1.11, based

			•		
Species	N _F	N _P	N _L	Ĥ	H range
Y. evonymellus	1	3	41.3	0.1380 ± 0.0123	0.1270-0.1513
Y. cagnagellus	1	7	29.0	0.1253 ± 0.0196	0.1004-0.1523
Y. mahalebellus	1	3	41.7	0.0587 ± 0.0167	0.0444-0.0771
Y. malinellus	2	6	34.7	0.0809 ± 0.0095	0.0704-0.0888
Y. padellus	>10	9	30.1	0.1272 ± 0.0149	0.1135-0.1554

Table 4. Heterozygosity levels (H) in five species closely related to Y. rorellus. N_F = number of food plants; N_P = number of populations studied; N_L = mean number of loci studied per species.

on 1,214 individuals). The general fit with Hardy-Weinberg proportions indicates an amphimictic breeding system as well.

A severe bottleneck when Y. rorellus arose may have caused the drop in the level of variation (Nei et al., 1975; Chakraborty and Nei, 1977). After a population reduction a bimodality in h levels of different loci can be observed (Nei et al., 1975; Berlocher, 1976). This has not been the case in Y. rorellus. It seems that the founders did not carry over genetic variation in the form of individual heterozygosity. Most of the variation found in Y. rorellus is unique (Table 3) and probably resulted from mutations after its original bottleneck. After a bottleneck, it takes more than $2N_e/(4N_e v + 1)$ generations (with N_e = effective population size and v = mutation rate/generation/locus) for the original H level to be restored.

Reliable estimates of population sizes in Yponomeuta are not available, let alone effective population sizes. Locally, Y. rorellus numbers can exceed 105, but population size is known to fluctuate heavily over years (Herrebout et al., 1976; A. M. Emmet, pers. comm.). Moreover, population differentiation and substructuring are difficult to assess, as the low level of variation does not allow the calculation of F_{ST} values (Wright, 1951), except in the case of G3pdh. The mean $F_{\rm ST}$ value over three years (0.0304 \pm 0.0304) indicates little differentiation among populations, although there is a striking difference in H between populations on S. alba and those on other *Salix* species (Table 1).

The time since Y. rorellus shared a common ancestor with its closest relative can be estimated from the genetic distance (Nei, 1971) between them to be 1.4×10^6 to 5.7×10^6 , depending on the formula used (Thorpe, 1982; Menken, 1982a). Bottlenecks lead to overestimates of divergence times (Chakraborty and Nei, 1977); the higher estimate

is probably the more reliable one (Thorpe, 1982). Because Y. rorellus is univoltine, there have thus been several million generations for H to have been restored. Using the crude estimates $N_{\rm e} = 10^5$ and $v = 10^{-7}$, the number of generations required is about 1.92 × 10⁵. A single bottleneck when Y. rorellus arose cannot, therefore, explain sufficiently its low H value. The difficulty of estimating the crucial quantities of D, N_e , v, and t means that it is hard to choose among various possible explanations, but some remarks can still be made. Thus, recurrent populationsize fluctuations (which do occur frequently in Y. rorellus) may have kept effective population sizes small, preventing populations from building up H. The absence of differential allelic fixation counts against this scenario though: it is possible that the present populations derived from a single recent population of restricted size, a bottleneck which may have been caused by a recent ice age (some 10,000 years ago). However, congeners probably experienced similar bottlenecks as they feed on plants belonging to the same plant community (Herrebout et al., 1976). The finding that Y. rorellus populations from Darmstadt and Bayreuth (West Germany) show the same dearth of variation (Menken, unpubl.), together with the extraordinarily simple sex-pheromone (Löfstedt et al., 1986) and the reduced chromosome number (n = 29) (Thorpe, 1929; Gershenson, 1967) of Y. rorellus compared with other *Yponomeuta* species (n = 31), all hint at a bottleneck at the origin of this species.

Other factors, both genetic and environmental, may reinforce the low incidence of new variation in *Y. rorellus*.

The number of food-plant species is an important component of niche breadth (environmental heterogeneity). A positive correlation between number of food plants and

genetic variation has been found for Rhagoletis (Berlocher, 1976), but not for a number of forest lepidopterous species (Mitter and Futuyma, 1980). In related Yponomeuta species, no clear correlation between H level and number of food plants could be found. Y. rorellus infesting various Salix species has by far the lowest H level; Y. cagnagellus and Y. evonymellus (both strictly monophagous), and Y. padellus (oligophagous) have H levels between 10 and 15% (Table 4). The importance of this and of factors such as low mutation rates, trophic stability (Levins, 1968), and effective population size is difficult to assess in the absence of critical data. They all remain in the realm of conjecture.

Yponomeuta species can be hybridized easily in the laboratory, but the monomorphic nature and genetic homogeneity of natural populations of Y. rorellus provide additional evidence of the effectiveness of its reproductive isolation from sympatric relatives in the padellus complex (Menken, 1980b).

Acknowledgments

Grateful acknowledgment is given to Guy Bush, Todd Bierbaum, Dan Howard, Paola Arduino, and Wim Herrebout for helpful criticisms on the manuscript and to Karla ter Horst and Stephanie Gauthier for typing the manuscript. This investigation was supported by the Netherlands Organization for the Advancement of Pure Research (ZWO/BION).

LITERATURE CITED

- Archie, J. W. 1985. Statistical analysis of heterozygosity data: Independent sample comparisons. Evolution 39:623-637.
- Ayala, F. J. 1982. Genetic variation in natural populations: Problem of electrophoretically cryptic alleles. Proc. Nat. Acad. Sci. USA 79:550-554.
- Berlocher, S. H. 1976. The genetics of speciation in *Rhagoletis* (Diptera: Tephritidae). Ph.D. Diss. Univ. Texas, Austin.
- CHAKRABORTY, R., AND M. NEI. 1977. Bottleneck effects on average heterogosity and genetic distance with the stepwise mutation model. Evolution 31: 347–356.
- FUTUYMA, D. J. 1979. Evolutionary Biology. Sinauer, Sunderland, MA.
- Gershenson, Z. S. 1967. Karyotype of the willow ermine moth *Yponomeuta rorellus* Hb. (Lepidoptera, Yponomeutidae). Tsitol. Genet. 1:79–80 (In Russian).

- Graur, D. 1985. Gene diversity in Hymenoptera. Evolution 39:190-199.
- HERREBOUT, W. M., P. J. KUIJTEN, AND J. T. WIEBES. 1976. Small ermine moths of the genus *Yponomeuta* and their host relationships (Lepidoptera, Yponomeutidae). Symp. Biol. Hung. 16:91–94.
- KIMURA, M. 1982. The neutral theory as a basis for understanding the mechanism of evolution and variation at the molecular level, pp. 3-56. In M. Kimura (ed.), Molecular Evolution, Protein Polymorphism and the Neutral Theory. Japanese Scientific Press, Tokyo, Japan.
- KIMURA, M., AND J. F. CROW. 1964. The number of alleles that can be maintained in a finite population. Genetics 49:725–738.
- KING, J. L., AND T. OHTA. 1975. Polyallelic mutational equilibria. Genetics 79:681–691.
- LEVINS, R. 1968. Evolution in Changing Environments. Princeton Univ. Press, Princeton, NJ.
- LÖFSTEDT, C., W. M. HERREBOUT, AND J.-W. DU. 1986. Filling of an ecological vacancy: Evolution of tetradecyl acetate as an ermine moth pheromone. Nature 323:621-623.
- Menken, S. B. J. 1980a. Inheritance of allozymes in *Yponomeuta I*. Intraspecific crosses in *Y. cagnagellus*. Proc. Kon. Ned. Acad. Wet. (C) 83:417-423.
- ——. 1980b. Inheritance of allozymes in *Yponomeuta* II. Interspecific crosses within the *padellus*-complex and reproductive isolation. Proc. Kon. Ned. Acad. Wet. (C) 83:425–431.
- ——. 1980c. Allozyme polymorphism and the speciation process in small ermine moths (Lepidoptera, Yponomeutidae). Thesis Univ. Leiden, Leiden, Neth.
- —. 1981. Host races and sympatric speciation in small ermine moths, Yponomeutidae. Entomol. Exp. Appl. 30:280–292.
- 1982a. Biochemical genetics and systematics of small ermine moths. Z. Zool. Syst. Evol.-forsch. 20:131–143.
- ——. 1982b. Enzymatic characterization of nine endoparasite species of small ermine moths (Yponomeutidae). Experientia 38:1461–1462.
- Menken, S. B. J., and S. A. Ulenberg. 1983. Diagnosis of the agromyzids *Liriomyza bryoniae* and *L. trifolii* by means of starch gel electrophoresis. Entomol. Exp. Appl. 34:205–208.
- ——. 1986. Allozymatic diagnosis of four economically important *Liriomyza* species (Diptera, Agromyzidae). Ann. Appl. Biol. 109:41–47.
- MITTER, C., AND D. J. FUTUYMA. 1980. Population genetic consequences of feeding habits in some forest Lepidoptera. Genetics 92:1005–1021.
- Nei, M. 1971. Interspecific gene differences and evolutionary time estimated from electrophoretic data on protein identity. Amer. Natur. 105:385-398.
- NEI, M., T. MARUYAMA, AND R. CHAKRABORTY. 1975. The bottleneck effect and genetic variability in populations. Evolution 29:1–10.
- NEI, M., AND A. K. ROYCHOUDHURY. 1974. Sampling variances of heterozygosity and genetic distance. Genetics 76:379–390.
- Nevo, E., A. Beiles, and R. Ben-Shlomo. 1984. The evolutionary significance of genetic diversity: Ecological, demographic and life history correlates, pp. 13–213. *In* G. S. Mani (ed.), Evolutionary Dynam-

- ics of Genetic Diversity. Lecture Notes in Biomathematics, Vol. 53. Springer-Verlag, Berlin, W. Ger. Powell, J. R. 1975. Protein variation in natural populations of animals. Evol. Biol. 8:79–119.
- SIMON, C., AND J. ARCHIE. 1985. An empirical demonstration of the lability of heterozygosity estimates. Evolution 39:463–467.
- THORPE, J. P. 1982. The molecular clock hypothesis: Biochemical evolution, genetic differentiation and systematics. Ann. Rev. Ecol. Syst. 13:139–168.
- THORPE, W. H. 1929. Biological races in *Hyponomeuta padella* L. J. Linn. Soc. Zool. 36:621–634. VARVIO-AHO, S.-L. 1981. The effects of ecological
- differences on the amount of enzyme gene variation in Finnish water-strider (*Gerris*) species. Hereditas 94:35–39
- WILKINSON, C., G. BRYAN, S. B. J. MENKEN, AND E. J. VAN NIEUKERKEN. 1983. A clarification of the status of four taxa in the *Ectoedemia angulifasciella* group (Nepticulidae:Lepidoptera). Neth. J. Zool. 33: 211–224.
- WRIGHT, S. 1951. The genetical structure of populations. Ann. Eugen. 15:323-354.

Corresponding Editor: R. H. Crozier