

Absence of $M3$ Quenching in ^{26}Mg
[Phys. Rev. Lett. 74, 642 (1995)]

Kamal K. Seth, R. Soundranayagam, A. Saha, C. W. de Jager, H. de Vries, B. A. Brown, and B. H. Wildenthal

Equation (4) should be

$$\frac{\sum B(M3)_{\text{expt.}}}{\sum B(M3)_{\text{theo.}}} = \frac{49.8(52)}{48.3} = 1.03(11). \quad (4)$$

In addition, the Fig. 2 caption should read

FIG. 2. Comparison between experimental results and shell-model predictions for $\sum B(M3)$. For states above 8.4 MeV the level numbers do not indicate any claimed correspondence between the experimental and theoretical levels; they are only for bookkeeping purposes.

Table I (with its caption) should read as follows:

TABLE I. Excitation energies and $B(M3)$ values for the 3^+ states in ^{26}Mg .

State	Expt	$E(\text{keV})$ Theory	$B(M3) (10^2 \mu_N^2 \text{ fm}^4)$	
			Expt	Theory
3_1^+	3941	3921	3.4(11)	1.4
3_2^+	4350	4510	7.0(30)	4.5
3_3^+	6125	6268	1.9(4)	6.0
3_4^+	7242	7281	11.7(20)	9.8
3_5^+	7724	7602	2.3(7)	0.6
3_6^+	8248	8004	11.8(21)	9.7
3_7^+	8456	8404	4.0(13)	16.3
3_8^+	9423	9115	4.0(15)	0.07
3_9^+	9902	9304	1.6(15)	0.03
3_{10}^+	10213	9576	2.1(11)	0.01
$\sum B(M3)$			49.8(52)	48.4