Downloaded from UvA-DARE, the institutional repository of the University of Amsterdam (UvA)
http://dare.uva.nl/document/58466

File ID 58466
Filename LIST OF FIGURES

SOURCE (OR PART OF THE FOLLOWING SOURCE):
Type Dissertation
Title Community structure and habitat associations of lowland grassland birds in Nepal.
Author H.S. Baral
Faculty Faculty of Science
Year 2001
Pages 235

FULL BIBLIOGRAPHIC DETAILS:
http://dare.uva.nl/record/97445

Copyright

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other then for strictly personal, individual use.

UvA-DARE is a service provided by the library of the University of Amsterdam (http://dare.uva.nl)

LIST OF FIGURES

Figure 1.1. Proposed former distribution of riverine tall grasslands in southern Nepal, northern India and Bangladesh (adapted from Bell and Oliver 1992). Remaining tall grassland is almost entirely restricted to the protected areas indicated. Grassland would have occurred as an intermittent tract rather than continuous cover.

Figure 1.2. Present distribution of tall grasslands in lowland Nepal. All areas shown are protected areas. Starting from west 1: Sukila Phanta, 2: Bardia, 3: Chitwan and 4: Parsa and 5: Koshi Tappu in the east. Dark shaded areas indicate grassland habitat.

Figure 2.1. Diurnal variation in bird activities.
(transect 2, Chitwan during 1996-1998)

Map 2.1. Sukila Phanta Wildlife Reserve with transects shown.

Map 2.2. Royal Chitwan National Park with transects shown.

Figure 4.1. Accretion of the bird species richness over different sampling intensities (1, 3, 4, 5, 10 visits).

Figure 4.2. Accretion of the bird abundance over different sampling intensities (1, 3, 4, 5, 10 visits).

Figure 4.3. Mean (with SD) percentage accretion of bird species in grassland habitats on consecutive sampling units. Grassland habitats defined by distance from forest edge (all sites, forest sites, intermediate sites, open grassland sites).

Figure 4.4. Mean (with SE) accretion of bird species in grassland types across 105 sites.

Figures 4.5-4.13: Accretion of bird species at individual sites in different area and seasons.

Figure 4.14. Percentage accretion of species over DCA axis 1, species richness and log abundance. The proportion of species surveyed after 5 visits compared to 10 visits.

Figure 4.15. Percentage accretion of species over log proportion of high emergent vegetation, distance from forest and grass height. The proportion of species surveyed after 5 visits compared to 10 visits.

Figure 5.1. DCA axis scores plotted on 2 principal axes for grassland types classified by distance to forest across 412 sites in lowland grasslands of Nepal.

Figures 5.2. DCA axis scores plotted on the principal axes for grassland types classified by grass assemblage across 412 sites in lowland grasslands of Nepal.

Figures 5.3. Centroids (with SD) on the principal DCA axes produced from the ordination of sites classified by grass assemblage and distance to forest.

Figure 5.4. DCA axis scores plotted on the principal axes for grassland types classified by distance to forests for winter and summer sites in lowland grasslands of Nepal.

Figure 5.5. Centroids (with SD) on the principal DCA axes for grassland types classified by distance to forests for winter and summer sites in lowland grasslands of Nepal.

Figure 5.6. DCA axis scores plotted on the principal axes for grassland types classified by grassland assemblage for winter and summer sites in lowland grasslands of Nepal.

Figure 5.7. Centroids (with SD) on the principal DCA axes for grassland types classified by grassland assemblage for winter and summer sites in lowland grasslands of Nepal.

Figure 5.8. Mean (with SD) on the principal DCA axes produced from the ordination of grassland bird assemblages for sites classified by grass assemblage and by distance to forests. The deviation in the DCA mean scores for two seasons are also indicated.

Figure 5.9. Dendrogram showing the 10 TWINSPAN groups in 176 sites visited in winter. The number of sites in each group is indicated below the TWINSPAN group number.

Figure 5.10. Dendrogram showing the 11 TWINSPAN groups in 236 sites visited in summer. The number of sites in each group is indicated below the TWINSPAN group number.

Figure 5.11. Pie charts showing the 10 TWINSPAN groups in 176 sites visited in winter showing the grassland habitat structure represented in them.

Figure 5.12. Pie charts showing the 11 TWINSPAN groups in 236 sites visited in summer showing the grassland habitat structure represented in them.
Figure 5.13. Pie charts showing the 10 TWINSPAN groups in 176 sites visited in winter and the 11 TWINSPAN groups in 236 sites visited in summer showing the grassland assemblages classified by grass species represented in them.
Figure 5.14. Mean (with SE) of grassland area, forest distance, grass height and emergent vegetations for 10 winter and 11 summer TWINSPAN groups.
Figure 5.15. Mean (with SE) of wild grazing, cattle grazing, cutting and burning for 10 winter and 11 summer TWINSPAN groups.
Figure 5.16. Mean (with SE) of species richness, abundance, threatened species richness and migrants (%) for 10 winter and 11 summer TWINSPAN groups.
Figure 5.17. Mean (with SE) of winter migrants (%) for 11 summer TWINSPAN groups. Note an exceptionally high number of winter migrants using TWINSPAN 6 group (wet Phragmites grassland).
Figure 5.18. Mean (with SE) of a) species richness b) grass height and c) DCA axis 3 for cattle grazing intensity at 46 Saccharum sites.
Figure 5.19. Dendrogram showing the 10 TWINSPAN groups in 176 sites visited in winter. The indicator bird species in each group are mentioned with their scores.
Figure 5.20. Dendrogram showing the 11 TWINSPAN groups in 236 sites visited in summer. The indicator bird species in each group are mentioned with their scores.
Figure 5.21. Grassland types and TWINSPAN groups represented in two study areas.
Figure 5.22. Frequency of occurrence of some bird species across winter TWINSPAN groups.
Figure 5.23. Frequency of occurrence of some bird species across summer TWINSPAN groups.
Figure 5.24. Frequency of occurrence of threatened bird species across winter TWINSPAN groups.
Figure 5.25. Frequency of occurrence of threatened bird species across summer TWINSPAN groups.
Figure 6.1. Percentage frequency of 60 most common species in winter and summer sites at Sukila Phanta Wildlife Reserve, Nepal.
Figure 6.2. Percentage frequency of 60 most common species in winter and summer sites at Royal Chitwan National Park, Nepal.
Figure 6.3. Mean (with SE) biological variables across various grassland types at 412 sites, lowland Nepal.
Figure 6.4. Mean (with SE) biological variables across various grassland types in winter and summer season, lowland Nepal
Figure 6.5. The percentage frequency of occurrence of threatened bird species in grassland habitats classified by their distance from forest (DF) and proportion of emergent vegetation. Forest edge grasslands (DF < 100m), intermediate grasslands (DF 100-500m), open grasslands (DF > 500m).
Figure 6.6. The percentage frequency of occurrence of threatened bird species in grassland habitats classified by dominant grass species.
Figure 6.7. The percentage frequency of occurrence of threatened bird species in lowland grassland during winter and summer seasons in two protected areas of Nepal. Status; r = resident; w = winter migrant; s = summer migrant; pm = passing migrant.
Figure 6.8. CCA of grassland bird communities in Royal Chitwan National Park
Figure 6.9. CCA of grassland bird communities, Sukila Phanta Wildlife Reserve
Figure 6.10. The percentage frequency of occurrence of bird species in grassland habitats classified by their distance to forest (DF) and proportion of emergent vegetation. Forest edge grasslands (DF<100m), intermediate grasslands (DF 100-500m), open grasslands (DF >500m).
Figure 6.11. The percentage frequency of occurrence of bird species in grassland habitats classified by dominant grass species.
Figure 6.12. The percentage frequency of occurrence of 25 most frequent bird species in grassland habitats classified by their distance to forest (DF) and proportion of emergent vegetation. Forest

edge grasslands (DF<100m), intermediate grasslands (DF 100-500m) with >10% emergent vegetation and <10% emergent vegetation and open grasslands (DF >500m).

Figure 6.13. The percentage frequency of occurrence of 25 most frequent bird species in grassland habitats classified by dominant grass species.

Figure 6.14. The percentage frequency of occurrence of bird species in grasslands of two lowland protected areas of Nepal during winter and summer seasons. Status; r = resident, w = winter migrant, s = summer migrant, pm = passage migrant.

Figure 7.1. Mean (with SE) soil moisture, cattle grazing, cutting, grass height and burning for grassland habitat structure classified by distance to forests during winter and summer seasons.

Figure 7.2. Mean (with SE) of soil moisture, cattle grazing, cutting, grass height and burning for different grassland assemblages as classified by grass species composition during winter and summer seasons.

Figure 7.3. Mean (with SE) of grass height, cutting, cattle grazing and burning for two different study sites during winter and summer seasons.

Figure 7.4. Mean (with SE) of grassland area, forest distance, grass height, soil moisture, water distance and emergent vegetation across all sites and season in grasslands classified by distance to forest.

Figure 7.5. Mean (with SE) of grassland area, forest distance, grass height, soil moisture, water distance and emergent vegetation across all sites and season.

Figure 7.6. Mean (with SE) of grass height and species richness in grasslands with different intensity of cattle grazing.