Downloaded from UvA-DARE, the institutional repository of the University of Amsterdam (UvA) http://dare.uva.nl/document/72877

File ID 72877 Filename Contents

SOURCE (OR PART OF THE FOLLOWING SOURCE):

Type Dissertation

Title Chemokine receptor signals: role in migration, invasion and cancer

metastasis

Author I.S. Zeelenberg Faculty Faculty of Medicine

Year 2004 Pages 135

FULL BIBLIOGRAPHIC DETAILS:

http://dare.uva.nl/record/203016

Copyright

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other then for strictly personal, individual use.

Contents

Chapter 1	Introduction	9	
Chapter 2	Retention of CXCR4 in the endoplasmic reticulum blocks dissemination of a T cell hybridoma. J. Clin. Invest. 108:269-277 (2001)		
Chapter 3	Stromal cell-derived factor-1-induced LFA-1 activation during in vivo migration of T cell hybridoma cells requires Gq/11, RhoA, and myosin, as well as Gi and Cdc42. J. Immunol. 166(7):4293-4301 (2001)		
Chapter 4	Blocking G-protein βγ dimers reduces desensitization of CXCR4, causing enhanced and persistent migration, independent of PI3-kinase. Submitted for publication	73	
Chapter 5	Fusion of docked vesicles, driven by synaptotagmin, is required for chemokine-induced T-cell migration. To be submitted		
Chapter 6	The chemokine receptor CXCR4 is required for outgrowth of colon carcinoma micrometastases. Cancer Res. 63:3833-3839 (2003)		
Chapter 7	A putative role for the chemokine receptor CXCR5 in tumor growth of colon carcinoma cells. Preliminary report		
Summary		117	
Samenvatting		121	
Dankwoord		127	
Curriculum vitae		133	
Liet of publications		135	