

File ID 98448
Filename Chapter 6 Summary and discussion

SOURCE (OR PART OF THE FOLLOWING SOURCE):

Type Dissertation
Title Identification and functional analysis of genes regulated by β 1 and β 3 integrins
Author J.I. van den Bout
Faculty Faculty of Medicine
Year 2008
Pages 112

FULL BIBLIOGRAPHIC DETAILS:

<http://dare.uva.nl/record/268974>

Copyright

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use.

Chapter 6:

SUMMARY AND DISCUSSION

Summary and Discussion

Integrins are heterodimeric transmembrane proteins that adhere to the extracellular matrix to mediate cell adhesion [1]. With their cytoplasmic tails integrins can bind to numerous proteins that link the integrin to the actin cytoskeleton and will initiate downstream signaling pathways as reviewed in **chapter 1**. These signaling pathways are involved in the regulation a variety of processes including migration, proliferation and differentiation [2].

While a large body of literature has been published on the effects of integrins on downstream effector proteins, less is known about the signals originating from the integrins that control gene transcription. There are several examples of genes regulated by integrins such as *cdc2* whose expression is increased when $\beta 3$ is reconstituted in a $\beta 3$ knockout cell line [3] and uPAR whose expression is decreased after overexpression of $\beta 3$ in CHO cell [4]. Moreover, when integrins interact with their ligand such as $\alpha 5\beta 1$ with fibronectin or $\alpha v\beta 3$ with vitronectin Bcl-2 expression is increased [5]. This thesis describes the identification of several genes whose transcription is regulated by integrins.

In this thesis we made extensive use of the GE11 cell line. The GE11 cell line is a neuro epithelial cell line isolated from the embryo of a $\beta 1$ knockout mouse. Subsequently, human $\beta 1$ has been reconstituted resulting in GE $\beta 1$ cells or human $\beta 3$ has been overexpressed resulting in GE $\beta 3$ [6], [7]. Expression of $\beta 1$ results in the loss of cell-cell contacts and the cells change from the epithelial-like morphology to become fibroblast-like. Additionally, these cells adhere better to fibronectin and migrate faster than the GE11 cells. On the other hand, $\beta 3$ overexpression also leads to a loss of cell-cell contacts but the cells become flat and well spread with many focal contacts present over the whole basal membrane of the cell. The difference between these two cell lines has been investigated previously and it was shown that in cells expressing $\beta 1$ the activity of the small GTPase RhoA is much higher than in cells in which $\beta 3$ is overexpressed [7].

In **chapter 2** we analyzed the effect on gene transcription by the reconstitution of $\beta 1$. Micro array analysis of the two cell lines GE11 and GE $\beta 1$ was used to identify genes whose transcription levels differed. Three genes were identified whose transcription levels were altered: that of the genes coding for thymosin- $\beta 10$ and insulin-like growth factor binding protein 4 (IGFBP-4) were decreased and that of the gene coding for galectin-3 was strongly increased. Galectin-3 was chosen for further investigation because it can directly bind to integrins such as $\alpha 1\beta 1$ [8] and can also influence integrin-mediated adhesion [9]. Furthermore, it is a prognostic marker for several cancers [10]. The hypervariable region of the I-like domain of $\beta 1$ was shown to regulate the activity of RhoA [11] while recent observations show that it also regulates cell contractility [12]. In this study we show that galectin-3 expression is dependent on the presence of this region. However, we could not demonstrate that galectin-3 expression is dependent on the activity of RhoA. Therefore, we suggest that two separate pathways downstream of $\beta 1$ regulate RhoA activity and galectin-3 expression independently. Further studies show that the increased expression of galectin-3 is important for enhanced cell adhesion to fibronectin and increased migration. While this data concurs with other reports showing that overexpression of galectin-3 increases adhesion the mode of action galectin-3 employs to enhance adhesion still remains unclear. Interestingly, in a recent paper it was shown that galectin-3 is involved in integrin-independent adhesion to the extracellular matrix through binding to the N-glycan chains attached to the matrix proteins [13]. On the other hand, galectin-3 has also been shown to mediate integrin-dependent adhesion [14], [15].

Interestingly, our data shows that galectin-3 is not involved in the maintenance of the $\beta 1$ -mediated cell morphology or in fibronectin fibrillogenesis, both of which are regulated by RhoA activity. Since galectin-3 expression is also not regulated by RhoA activity we conclude that galectin-3 expression and RhoA activity are regulated by separate pathways and that both in turn regulate different characteristics of GE $\beta 1$ cells.

Not only the reconstitution of $\beta 1$ but also the overexpression of $\beta 3$ leads to dramatic changes in cell morphology and behavior such as the loss of cell-cell contacts, increased cell spreading, migration and the formation of numerous focal adhesions [7]. Since $\beta 3$ overexpression was reported to regulate uPAR expression in CHO cells [4] we investigated whether $\beta 3$ overexpression in GE11 cells influences gene expression. In **chapter 3** we describe the identification of the gene coding for MRP that is downregulated when $\beta 3$ is overexpressed. MRP is an appealing target of regulation by $\beta 3$ since it can bind to actin

filaments [16] and can regulate the diffusion of $\beta 3$ integrins on the cell surface [17], [18]. It has been shown that MRP expression can be regulated by stimuli such as LPS [19] while the expression of the close relative of MRP, MARCKS, is regulated by transformation with v-Src or p21-Ras [20]. Our investigation shows that the Ras/MAPK pathway regulates the basal expression of MRP in the GE11 cell line but $\beta 3$ bypasses this pathway to downregulate MRP expression. The expression of deletion mutants of $\beta 3$ showed that the last four amino acids of the cytoplasmic tail of $\beta 3$ were important for MRP expression. Since this region of $\beta 3$ was also shown to bind to Src [21] we tested if Src controlled MRP expression. However, this was not the case. Therefore, we conclude that a Src-independent pathway involving the binding of another protein to this region could be important for the regulation of MRP expression. Interestingly, this same region of $\beta 3$ was also important for the increased cell spreading seen in GE $\beta 3$ cells. However, knockdown of MRP in GE11 or overexpression of MRP in GE $\beta 3$ cells had no effects on cell spreading, focal adhesion formation or cell morphology suggesting that MRP expression alone does not able to influence the morphology of the cells.

We continued our investigation of MRP to try to understand the mechanism MRP employs to regulate its localization within the cell. MRP uses a myristoylation motif and a polybasic cluster called the effector domain to bind to the plasma membrane in a similar way as a large group of proteins including Src, K-Ras and MARCKS. In **chapter 4** we describe how MRP localization is regulated. Controversy exists about the question how these proteins are specifically localized at the plasma membrane and not at the internal membranes that make up a larger surface area. Although it is generally accepted that these proteins are dependent on acidic phospholipids for plasma membrane targeting, it is still unknown which specific phospholipids are important. Recently, it was shown that proteins such as K-Ras use PIP₂ and PIP₃ to target to the plasma membrane [22] while a contrasting study showed that K-Ras targets to the plasma membrane because of the presence of the monovalent acidic lipid, phosphatidyl serine [23]. We show that loss of PIP₂ and PIP₃ does not affect the targeting or the association of MRP to the plasma membrane. Therefore, we suggest that phosphatidyl serine is responsible for the targeting of MRP. Furthermore, we tested if the myristoyl-electrostatic switch model applies to the regulation of MRP localization as it does for MARCKS. This model predicts that the combination of the binding of the myristoyl moiety and the positively charged effector domain are needed for the association of the protein with the membrane [24]. From our data we can conclude that this model applies for the association of MRP with the plasma membrane.

MRP associates with the plasma membrane but upon phosphorylation of the serine residues within the effector domain the protein is translocated away from the plasma membrane to internal membranes such as those of the lysosomes in a matter of seconds. Unlike MARCKS whose translocation from the plasma membrane to the internal membranes only occurred after several hours [25] MRP was so rapidly translocated to internal membranes that it was not possible to detect an increase in cytoplasmic MRP using fractionation assays. When the effector domain was deleted from the protein, MRP was no longer able to bind to the plasma membrane but was still present on the internal membranes. This raises the question how MRP associates with these internal membranes under these circumstances. If the myristoyl-electrostatic switch model also applies to the binding of MRP to internal membranes it would be impossible for MRP to bind when the effector domain is deleted. Therefore, the model explains the association of MRP to the plasma membrane through the cumulative effect of the myristoyl moiety and the effector domain binding to the plasma membrane but we suggest that MRP only utilizes the myristoyl moiety for association with internal membranes. Thus, the effector domain functions primarily as a plasma membrane targeting motif.

In different cancers an increase in c-Src activity has been observed even though activating mutations in the *SRC* gene are rare. While this argues against a role for c-Src in tumor initiation, it has been implicated in the formation of metastases and tumor growth. In **chapter 5**, it is shown, that the expression of “primed” c-Src carrying a mutation at the tyrosine 530 that disrupts intramolecular association between this residue and the Src homology domain 2 (SH2), in cells expressing $\beta 3$, but not $\beta 1$, promotes tumorigenicity. The cooperation between $\alpha v\beta 3$ and c-Src seems to be important since increased $\alpha v\beta 3$ expression has been linked to growth and tumor progression in cancers in which c-Src activity is often increased. In this chapter we show that $\alpha v\beta 3$ and primed c-Src are functionally linked to promote tumorigenicity in the cell lines tested. Others showed that there is a direct association between the SH3 domain of Src and the C-terminal YRGT motif of $\beta 3$ [21]. We found that the presence of this YRGT motif of $\beta 3$ is essential for the increased tumorigenicity.

Moreover, Src activity is also increased in cells expressing $\beta 3$. Src is activated in a stepwise process. Clustering of $\alpha v\beta 3$ brings the associated primed Src molecules in close contact with one another allowing their autophosphorylation in trans resulting in their full activation. Therefore, we conclude that the binding of Src to $\beta 3$ allows the clustering of Src resulting in the activation of Src leading to increased cell growth and tumorigenicity.

To conclude, we have identified several new targets of regulation by integrins and investigated their role in the integrin-mediated effects observed in our cell system. Furthermore, we have shed light on the mechanism by which MRP is targeted to the plasma membrane and how its association with this membrane is regulated. Lastly, we have shown that there is a functional link between Src and $\beta 3$ during tumorigenesis.

References

1. J.W.Tamkun, D.W.DeSimone, D.Fonda, R.S.Patel, C.Buck, A.F.Horwitz, R.O.Hynes (1986). Structure of integrin, a glycoprotein involved in the transmembrane linkage between fibronectin and actin. *Cell* **46**, 271-282.
2. R.O.Hynes (2002). Integrins: bidirectional, allosteric signaling machines. *Cell* **110**, 673-687.
3. T.Manes, D.Q.Zheng, S.Tognin, A.S.Woodard, P.C.Marchisio, L.R.Languino (2003). Alpha(v)beta3 integrin expression up-regulates cdc2, which modulates cell migration. *J Cell Biol* **161**, 817-826.
4. S.Hapke, M.Gawaz, K.Dehne, J.Kohler, J.F.Marshall, H.Graeff, M.Schmitt, U.Reuning, E.Lengyel (2001). beta(3)A-integrin downregulates the urokinase-type plasminogen activator receptor (u-PAR) through a PEA3/ets transcriptional silencing element in the u-PAR promoter. *Mol Cell Biol* **21**, 2118-2132.
5. M.L.Matter, E.Ruoslahti (2001). A signaling pathway from the alpha5beta1 and alpha(v)beta3 integrins that elevates bcl-2 transcription. *J Biol Chem* **276**, 27757-27763.
6. C.Gimond, A.Der Flier, S.van Delft, C.Brakebusch, I.Kuikman, J.G.Collard, R.Fassler, A.Sonnenberg (1999). Induction of cell scattering by expression of beta1 integrins in beta1-deficient epithelial cells requires activation of members of the rho family of GTPases and downregulation of cadherin and catenin function. *J Cell Biol* **147**, 1325-1340.
7. E.H.Danen, P.Sonneveld, C.Brakebusch, R.Fassler, A.Sonnenberg (2002). The fibronectin-binding integrins alpha5beta1 and alphavbeta3 differentially modulate RhoA-GTP loading, organization of cell matrix adhesions, and fibronectin fibrillogenesis. *J Cell Biol* **159**, 1071-1086.
8. J.Ochieng, M.L.Leite-Browning, P.Warfield (1998). Regulation of cellular adhesion to extracellular matrix proteins by galectin-3. *Biochem Biophys Res Commun* **246**, 788-791.
9. I.Kuwabara, F.T.Liu (1996). Galectin-3 promotes adhesion of human neutrophils to laminin. *J Immunol* **156**, 3939-3944.
10. S.Califice, V.Castronovo, F.Van Den Brule (2004). Galectin-3 and cancer (Review). *Int J Oncol* **25**, 983-992.
11. H.Miao, S.Li, Y.L.Hu, S.Yuan, Y.Zhao, B.P.Chen, W.Puzon-McLaughlin, T.Tarui, J.Y.Shyy, Y.Takada, S.Usami, S.Chien (2002). Differential regulation of Rho GTPases by beta1 and beta3 integrins: the role of an extracellular domain of integrin in intracellular signaling. *J Cell Sci* **115**, 2199-2206.
12. S.Huveneers, R.Fassler, A.Sonnenberg, E.H.Danen. Integrin 51 Couples High Affinity Fibronectin Binding to Rho-mediated Contractility and Fibrillogenesis. 2007. Ref Type: Unpublished Work
13. J.Friedrichs, J.M.Torkko, J.Helenius, T.P.Teravainen, J.Fullekrug, D.J.Muller, K.Simons, A.Manninen (2007). Contributions of galectin-3 and -9 to epithelial cell adhesion analysed by single cell force spectroscopy. *J Biol Chem*.
14. A.Lagana, J.G.Goetz, P.Cheung, A.Raz, J.W.Dennis, I.R.Nabi (2006). Galectin binding to Mgat5-modified N-glycans regulates fibronectin matrix remodeling in tumor cells. *Mol Cell Biol* **26**, 3181-3193.
15. P.Matarrese, O.Fusco, N.Tinari, C.Natoli, F.T.Liu, M.L.Semeraro, W.Malorni, S.Iacobelli (2000). Galectin-3 overexpression protects from apoptosis by improving cell adhesion properties. *Int J Cancer* **85**, 545-554.
16. F.Wohnsland, M.O.Steinmetz, U.Aebi, G.Vergeres (2000). MARCKS-related protein binds to actin without significantly affecting actin polymerization or network structure. Myristoylated alanine-rich C kinase substrate. *J Struct Biol* **131**, 217-224.
17. T.Jin, J.Li (2002). Dynamitin controls Beta 2 integrin avidity by modulating cytoskeletal constraint on integrin molecules. *J Biol Chem* **277**, 32963-32969.
18. X.Zhou, J.Li (2000). Macrophage-enriched myristoylated alanine-rich C kinase substrate and its phosphorylation is required for the phorbol ester-stimulated diffusion of beta 2 integrin molecules. *J Biol Chem* **275**, 20217-20222.
19. J.R.Sunohara, N.D.Ridgway, H.W.Cook, D.M.Byers (2001). Regulation of MARCKS and MARCKS-related protein expression in BV-2 microglial cells in response to lipopolysaccharide. *J Neurochem* **78**, 664-672.
20. J.C.Reed, U.Rapp, M.P.Cuddy (1991). Transformed 3T3 cells have reduced levels and altered subcellular distribution of the major PKC substrate protein MARCKS. *Cell Signal* **3**, 569-576.
21. E.G.Arias-Salgado, S.Lizano, S.Sarkar, J.S.Brugge, M.H.Ginsberg, S.J.Shattil (2003). Src kinase activation by direct interaction with the integrin beta cytoplasmic domain. *Proc Natl Acad Sci U S A* **100**, 13298-13302.

22. W.D.Heo, T.Inoue, W.S.Park, M.L.Kim, B.O.Park, T.J.Wandless, T.Meyer (2006). PI(3,4,5)P3 and PI(4,5)P2 lipids target proteins with polybasic clusters to the plasma membrane. *Science* **314**, 1458-1461.
23. T.Yeung, M.Terebiznik, L.Yu, J.Silvius, W.M.Abidi, M.Philips, T.Levine, A.Kapus, S.Grinstein (2006). Receptor activation alters inner surface potential during phagocytosis. *Science* **313**, 347-351.
24. S.McLaughlin, A.Aderem (1995). The myristoyl-electrostatic switch: a modulator of reversible protein-membrane interactions. *Trends Biochem Sci* **20**, 272-276.
25. L.A.Allen, A.Aderem (1995). Protein kinase C regulates MARCKS cycling between the plasma membrane and lysosomes in fibroblasts. *EMBO J* **14**, 1109-1121.